

8-15-2008

The Urysohn Space Embeds in Banach Spaces in Just One Way

M. Randall Holmes
Boise State University



This is an author-produced, peer-reviewed version of this article. © 2009, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>). The final, definitive version of this document can be found online at *Topology and its Applications*, doi:10.1016/j.topol.2008.03.013

The Urysohn Space Embeds in Banach Spaces in Just One Way

M. Randall Holmes

This note will give a brief account of the research behind the author's master's thesis of 1985 in which we gave an account of the Urysohn universal separable metric space, which we had discovered independently but which was of course not new (see [4]), and of the unique universal separable Banach space which appears as the linear closure of any isometric copy of the Urysohn space in a Banach space, of which we gave the first account (the complete reference on our work in this area is [2]).

The question which I was asked in a graduate general topology class was “Is there a universal separable metric space (implicitly, up to homeomorphism)”? That is, is there a metric space U such that for any separable metric space at all, there is a homeomorphic embedding from X into U ?

Now of course an isometry is a homeomorphism, and we perhaps foolishly asked the harder question “Is there a universal separable metric space up to isometry”?

To investigate this question, we defined the concept of a “possible combination of distances” from a metric space X .

Let (X, d) be a metric space (which we will refer to as X , as is usual (if logically dubious)). The metrics on all spaces will be d as long as the intended space can be understood from context: otherwise the metric on X will be d_X .

Definition 1. Let $Y \subseteq X$. Let p be a function from Y to the non-negative reals, satisfying $p(u) - p(v) \leq d(u, v) \leq p(u) + p(v)$ for all u and v in Y . (Of course $|p(u) - p(v)| \leq d(u, v)$ then holds by symmetry). Such a function will be called a *possible combination of distances from Y (as a subspace of X)*.

If the domain of p is X , we can adjoin a new point q to X , stipulating that $d(q, x) = p(x)$ for each $x \in X$, and it is straightforward to verify that $X \cup \{q\}$

is a metric space with this metric. If the domain Y of p is a proper subset of X , one can extend p to the whole of X by making its value at each $x \in X - Y$ as large as possible: let $p'(x)$ be defined as $\inf\{d(x, y) + p(y) \mid y \in Y\}$. It is straightforward to verify that p' agrees with p on Y and is a possible combination of distances from the whole of X . Thus it is possible to adjoin a point to X in such a way that its distance from every point y of Y is $p(y)$. The notion “possible combination of distances from Y ” exactly captures the possible combinations of distances from Y of a new point to be adjoined to the ambient metric space X .

Definition 2. Of course possible combinations of distances from a proper subset Y may be handled by points already found in $Y - X$: we say that $x \in X$ realizes p if $d(x, y) = p(y)$ for each $y \in Y$.

The universal separable metric space of Urysohn can be characterized using this notion. Up to isometry, U is the unique complete metric space which has the property that any possible combination of distances from a finite subset of U is realized in U (this follows readily from the standard characterizations of the Urysohn space).

U can be constructed using this notion as well. Let X be a metric space and let X' be the set of all possible combinations of distances from the whole of X . We put the metric $d(p, q) = \sup\{|p(x) - q(x)| \mid x \in X\}$ on X' . X' has a canonical subspace isometric to X (consisting of the possible combinations of distances p_x which take on the value zero at some point x of X). Moreover, for any $p \in P$, we have $d(p, p_x) = p(x)$. The space X' contains new points realizing every possible combination of distances from (the natural isometric copy of) X , with any two new points as close to one another as their distances from the natural embedded copy of X permit. Unfortunately, X' is not as a rule separable; so we define a subspace X'' of X' as the completion in X' of the set of extensions to all of X (as described above) of possible combinations of distances from finite subsets of X . X'' can be shown to be separable if X is separable. Now let X_0 be a one-point space and define X_{n+1} as X''_n for each natural number n . The completion of the direct limit of the X_i 's (using the natural isometric embedding of each space in the sequence into the next to construct the direct limit) is isomorphic to U . This was our original construction of a universal separable metric space up to isometry when we discovered this space independently in 1983.

No one at SUNY Binghamton had heard of the Urysohn space, but many people there knew of the well-known theorem of Banach and Mazur that

$C[0, 1]$ (the space of continuous functions from $[0, 1]$ to the reals with the sup metric) is a universal separable metric space up to isometry (and in fact a universal separable Banach space up to linear isometry) (see [1]). So the natural question in our mind was “how do U and $C[0, 1]$ embed into each other?”

It was immediately clear that the spaces are different. Consider the constant functions 1,2,3. A possible combination of distances from these points which cannot be realized in $C[0, 1]$ maps each of these points to 1. But 1 is the only point in $C[0, 1]$ which is at distance 1 from each of 2 and 3.

The perhaps valuable original contribution of my work on the Urysohn space from 1983 to 1985 is contained in the following series of observations.

Definition 3. We define a *possible combination of values* of a set of functions $F \subseteq C[0, 1]$ as a function p from F to the reals such that for any $f, g \in F$, we have $|p(f) - p(g)| \leq d(f, g)$. Further, we say that a real r *realizes* p iff for each $f \in F$, we have $f(r) = p(f)$. It should be clear that for any $r \in [0, 1]$, the function sending each $f \in F$ to $f(r)$ is a possible combination of values for F (justifying the terminology).

Suppose that F is a finite subset of an isometric copy of U in $C[0, 1]$, $0 \in F$, and p is a possible combination of values for F . It is straightforward to verify that for large enough N (twice the diameter of F will work) the function ($f \in F \mapsto N - p(f)$) is a possible combination of distances from F , and so there is a function g in the isometric copy of U which realizes these distances from F . Extend p by defining $p(g) = N$ (obviously the extended p is still a possible combination of values). Now further it is straightforward to show that ($f \in F \cup \{g\} \mapsto N + p(f)$) is a possible combination of distances from $F \cup \{g\}$, so there is a function h in the isometric copy of U which realizes these distances from $F \cup \{g\}$. Now $d(g, h) = 2N$ by construction, so there must be a real r_p such that $|g(r_p) - h(r_p)| = 2N$. Since $0 \in F$ and $d(g, 0) = d(h, 0) = N$, we are forced to have either $f(r_p) = N$ and $g(r_p) = -N$ or $f(r_p) = -N$ and $g(r_p) = N$. For each $f \in F$, we have $|f(r_p) - g(r_p)| \leq N - p(f)$ and $|f(r_p) - h(r_p)| \leq N + p(f)$. So in the first case $f(r_p)$ is forced to have the value $p(f)$ for each $f \in F$ and in the second case $f(r_p)$ is forced to have the value $-p(f)$ for each $f \in F$. So we have shown the following rather surprising

Theorem 4. *For any finite subset F of an isometric copy of U containing 0 with $0 \in F$, and any possible combination of values p for F , either p is realized at some $r_p \in [0, 1]$ or $-p$ is realized at some $r_p \in [0, 1]$.*

This is very strange! It implies for example, the following

Corollary 5. *Any element of an isometric copy of U in $C[0, 1]$ which contains 0, other than 0 itself, is a component of something which is almost a Peano curve.*

Proof. Let f be such a function. A possible combination of distances from f and 0 is the map sending 0 to $|f|$ and $|f|$ to $2|f|$, so there is a point f_2 at distance $|f|$ from 0 and $2|f|$ from f in the isometric copy of U . Any element of $[-|f|, |f|]^2$ is of the form $(p(f), p(f_2))$ where p is a possible combination of values for f and f_2 (and all possible p are associated with points in this way). Thus for every point (x, y) in $[-|f|, |f|]^2$ there is a real r such that either $f(r) = x$ and $f_2(r) = y$ or $f(r) = -x$ and $f_2(r) = -y$: f and f_2 are the components of a continuous curve which visits each point of a square centered at the origin or its mirror image through the origin. \square

So we see that no familiar function in $C[0, 1]$ except the constant 0 can be an element of such a copy of U ! What, on the face of it, does the universal separable metric space of Urysohn have to do with Peano curves?

Further, consider the linear closure of an isometric copy of U in $C[0, 1]$ containing 0. Consider in particular any finite linear combination $\sum c_i f_i$ of elements of the copy of U . The norm of $\sum c_i f_i$ is the supremum of all sums $|c_i f_i(r)|$ for $r \in [0, 1]$. But this means that it is the supremum of all sums $|c_i p(f_i)|$ where p is a possible combination of values for the set of f_i 's, because every such possible combination of values or its uniform negative is realized at some r . This supremum depends only on the distances among the f_i 's and 0, so such norms are determined entirely by the metric structure of U and the selection of a point to correspond to 0. This completes the proof of another surprising

Theorem 6. *The linear closure of an isometric copy of U in $C[0, 1]$ which contains 0 is a uniquely determined separable Banach space \overline{U} , up to linear isometry (and so, because of the known universality of $C[0, 1]$, the linear closure of an isometric copy of U containing 0 in any Banach space is uniquely determined up to linear isometry).*

There are two questions about this which present themselves. One of them was ours, on which we made little progress, but we were able to answer a question of Sierpinski.

Question 7. We know that U is a universal separable metric space up to isometry. Is its uniquely determined linear closure \overline{U} a universal separable Banach space up to linear isometry?

We did not make much headway on this. In [2] we got as far as demonstrating that \overline{U} did not have a certain homogeneity property which would have facilitated a proof of universality. We have been told in informal contexts that the answer to the question is now known to be positive but we have not received a satisfactory account of this yet.

The second question, which I did answer, is difficult to phrase precisely. The proof that $C[0, 1]$ is a universal separable Banach space under linear isometry involves space-filling curves. We present a version adapted to embedding metric spaces rather than Banach spaces (we believe this adaptation is from [3]). Let X be a metric space and fix an element of X which will be mapped to 0. Let D be a countable dense subset of X . Take the space D^* of all possible combinations of values of D (defined as above, but of course this was not their terminology) and put the pointwise convergence topology on it. This space is a connected compact metric space, so one can define a continuous map f from $[0, 1]$ onto D^* . Now with each point $d \in D$ associate the function which sends each $r \in [0, 1]$ to $f(r)(d)$. Under the supremum metric, these functions will make up an isometric copy of D in $C[0, 1]$ whose completion will be a copy of X . Sierpinski observed, in commenting on this proof in [3], that for most familiar spaces nothing as nasty as this construction using a Peano curve is required, and he asked specifically this

Question 8. (Sierpinski)

Is there a better way to embed U in $C[0, 1]$ than the general method of Banach and Mazur, as adapted to metric spaces?

The results above linking isometric embeddings of U with $C[0, 1]$ strongly suggest that the answer should be No. However, it is tricky to formulate the negative answer precisely.

In [2], we formulated precise conditions under which a finite subset F of $C[0, 1]$ can be extended to an isometric copy of U containing 0. The condition is equivalent to the statement that there is a positive constant N and a function g at distance $N + d(0, f)$ from f for each $f \in F$, such that for each possible combination of values p for $F \cup \{g\}$, either p is realized or $-p$ is realized. It follows easily from the discussion above that these conditions are necessary; additional work is required to show that these conditions are

sufficient. Such a set F is called *inflatable* in [2]. The basic idea of the proof is that one can choose any possible combination of distances p from F , then use g to guide the construction of two functions, a function f' which has the desired distances from the elements of f and a function g' which has distance $N + d(f, 0)$ from each $f \in F \cup \{f'\}$. This allows the construction of a countable dense subset of a copy of U from an inflatable set, and taking the completion of a subset of $C[0, 1]$ of course presents no difficulties. This allows an exposition of the construction of U entirely in terms of $C[0, 1]$, which is given in detail in [2].

An easy way to answer Sierpinski's question is the following: any embedding of U into $C[0, 1]$ is associated via the construction outlined above with a continuous curve in D^* (where D is a countable dense subset of U) which "half-fills" D^* (visits each element of D^* or its negative). So mod the difference between "half-space-filling curves" and frankly space-filling curves, the answer to the question of Sierpinski is indeed No. A more subtle approach involves choosing D cleverly so that a universal construction of isometric embeddings of U in $C[0, 1]$ can be presented whose only parameter is a "half-space-filling curve" in the usual Hilbert cube. This can be done in such a way that there is a one-to-one correspondence between half-space-filling curves and isometric embeddings, if D is chosen in such a way that all instances of the triangle inequality are strict (so no finite assignment of values at a given r to points of D can exactly fix the value at r of any other element of D) (it is noted that this can be done in [2] but complete details are not given for the more refined version).

References

- [1] S. Banach, *Théorie des Opérations Linéaires*, Hafner, New York 1932, p. 185.
- [2] M. Randall Holmes, "The universal separable metric space of Urysohn and isometric embeddings thereof in Banach spaces", *Fundamenta Mathematicae* vol 140 (1992), pp. 199-223.
- [3] W. Sierpiński, "Sur un espace métrique universel", *Fundamenta Mathematicae*, vol. 43 (1945), pp. 115-122. Also see his *General Topology*, Univ. of Toronto Press, Toronto 1952, pp. 159-162.

- [4] P. Urysohn, “Sur un espace metrique universel”, *Bull. Sci. Math.* 51 (1927), pp. 43-64, 74-90.