

BIOMECHANICAL EVALUATION OF GLENOHUMERAL JOINT STABILIZING  
MUSCLES DURING PROVOCATIVE TESTS DESIGNED TO DIAGNOSE  
SUPERIOR LABRUM ANTERIOR-POSTERIOR LESIONS

by

Vanessa J. C. Wood

A thesis

submitted in partial fulfillment

of the requirements for the degree of

Master of Science in Mechanical Engineering

Boise State University

Summer 2009

© 2009  
Vanessa J.C. Wood  
ALL RIGHTS RESERVED

BOISE STATE UNIVERSITY GRADUATE COLLEGE

**DEFENSE COMMITTEE AND FINAL READING APPROVALS**

of the thesis submitted by

Vanessa J.C. Wood

Thesis Title: Evaluation of Glenohumeral Joint Stabilizing Muscles During Provocative Tests Designed to Diagnose Superior Labrum Anterior-Posterior Lesions

Date of Final Oral Examination: 09 June 2009

The following individuals read and discussed the thesis submitted by student Vanessa J.C. Wood, and they evaluated her presentation and response to questions during the final oral examination. They found that the student passed the final oral examination.

Michelle B. Sabick, Ph.D. Chair, Supervisory Committee

Ron P. Pfeiffer, Ph.D. Member, Supervisory Committee

Kotaro Sasaki, Ph.D. Member, Supervisory Committee

The final reading approval of the thesis was granted by Michelle B. Sabick, Ph.D., Chair of the Supervisory Committee. The thesis was approved for the Graduate College by John R. Pelton, Ph.D., Dean of the Graduate College.

## ACKNOWLEDGEMENTS

At this time I would like to express my profound gratitude to a handful of people who were instrumental in my success at Boise State University and the completion of my degree. For the length of my studies at Boise State University, I have been extremely fortunate to receive extensive academic and financial support. My accomplishments would have been impossible without the guidance, patience, and generosity of these individuals.

- Thesis Committee
  - Dr. Michelle Sabick, Advisor
  - Dr. Ronald Pfeiffer
  - Dr. Kotaro Sasaki

I would also like to thank the following individuals for their help and support in the completion of this project specifically. Without them, this study would not have been possible.

- Center for Orthopaedic and Biomechanics Research
  - Seth Kuhlman
  - Jason Christensen
- Intermountain Orthopaedic
  - Dr. Michael Curtin
  - Dr. Kurt Nilsson
  - Dr. Kevin Shea

Furthermore I must express my gratitude for the opportunities I received here at Boise State as a Graduate Research Assistant for both Dr. Michelle B. Sabick (2007-2009) and Dr. John F. Gardner (2006-2007), without their support I clearly would not have been able to pursue graduate studies or academic research.

## ABSTRACT

### **Biomechanical Evaluation of Glenohumeral Joint Stabilizing Muscles during Provocative Tests Designed to Diagnose SLAP Lesions**

**Vanessa Wood**

Despite considerable advances in the understanding of glenohumeral (GH) biomechanics and glenoid labral pathologies, arthroscopy remains the only definitive means of Superior Labrum Anterior-Posterior (SLAP) lesion diagnosis. Unfortunately, natural GH anatomic variants limit the reliability of radiography. Accurate clinical diagnostic techniques would be advantageous due to the invasiveness, patient risk, and financial cost associated with arthroscopy. Twenty provocative tests designed to elicit labral symptoms as a diagnostic sign have shown promising accuracy by their respective original authors, but later studies generally fail to reproduce those findings. The purpose of this study was to compare the behavior of GH joint stabilizing muscles in promising tests. Electromyography (EMG) was used to characterize the activation of GH joint stabilizing muscles, with particular interest in the Long Head Biceps Brachii (LHBB) behavior, as activation of the LHBB and subsequent tension in the biceps tendon should illicit labral symptoms in SLAP lesion patients.

Volunteers (n=21) with no history of shoulder pathology were recruited for this study. The tests analyzed were Active Compression, Speed's, Pronated Load, Biceps Load I (Bicep I), Biceps Load II (Bicep II), Resisted Supination External Rotation (RSER), and Yergason's. Test modifications that allowed the use of the Biodex System improved reproducibility. EMG was used to record activity for GH muscles: the LHBB, short head of the biceps brachii, anterior deltoid, pectoralis major, latissimus dorsi, infraspinatus, and supraspinatus. An indwelling electrode was used to monitor supraspinatus activity, and the remaining muscles utilized surface electrodes. EMG data were recorded at 1250 Hz and filtered with custom MATLAB software. Muscle activity for each test was characterized by activation and selectivity. Muscle activation was defined as the muscle's peak normalized EMG amplitude. Muscle selectivity was defined as the ratio of muscle activation for the muscle of interest over the sum of all seven muscles' peak activations.

Results indicated that Bicep I and II had the greatest potential for the clinical detection of SLAP lesions because both tests 1) elicited large LHBB activation, suggesting that during these tests more tension was applied to the biceps tendon, and also 2) remained highly selective for the LHBB, which should reduce the potential sources for confounding results. Also, tests that elicited promising LHBB behavior for either a single suite or for both activation and selectivity, shared design patterns relating to location of the applied load, forearm orientation, joint position, and line of pull. These characteristics should be further examined to determine their potential role in optimizing SLAP test design and improving clinical diagnostic techniques.

## TABLE OF CONTENTS

ACKNOWLEDGEMENTS .....	v
ABSTRACT .....	vii
LIST OF TABLES .....	x
LIST OF FIGURES .....	xi
INTRODUCTION .....	1
Operational Definitions.....	7
Limitations .....	8
Delimitations.....	8
LITERATURE REVIEW .....	9
Shoulder Anatomy .....	9
SLAP Lesions .....	11
Provocative Tests .....	14
Active Compression Test.....	14
Speed's Test .....	17
Pronated Load Test .....	18
Bicep Load I Test.....	19
Bicep Load II Test .....	19
Resisted Supination External Rotation Test .....	20
Supination Sign Test.....	21
METHODS .....	22
Experimental Protocol .....	22
Subject and IRB approval.....	22

Electromyography Apparatus and Subject Preparation .....	22
Subject Protocol .....	24
Test Descriptions and Modifications .....	25
Active Compression Test (ACPU and ACPD) .....	26
Speed's Test .....	28
Pronated Load Test .....	29
Bicep Load I Test.....	29
Bicep Load II Test .....	30
Resisted Supination External Rotation Test .....	30
Supination Sign Test.....	31
Electromyography Analysis.....	32
Initial EMG Processing.....	32
EMG Filtering Technique .....	33
Normalizing Provocative Test Data with MVIC Maximums .....	33
Muscle Activation.....	33
Muscle Selectivity.....	33
Statistical Analysis.....	34
REFERENCES .....	35
APPENDIX .....	40
Manuscript for Submission to the Journal of Shoulder and Elbow Surgery	40

LIST OF TABLES


Table 1. MVIC Joint Positions and Resisted Maneuvers ..... 24

## LIST OF FIGURES

Figure 1: SLAP Lesion Injury Region .....	1
Figure 2: SLAP Lesion Injury Mechanism I.....	4
Figure 3: SLAP Lesion Injury Mechanism II .....	5
Figure 4: Shoulder Complex - Bony Anatomy .....	9
Figure 5: Glenohumeral Joint Anatomy .....	10
Figure 6: Electrode Placement for EMG .....	23
Figure 7: ACPU .....	27
Figure 8: ACPD .....	27
Figure 9: Speeds Starting Position.....	28
Figure 10: Bicep Load I.....	29
Figure 11: Bicep Load II.....	30
Figure 12: RSER Starting Position .....	31
Figure 13: Yergason's.....	32

## INTRODUCTION

The shoulder complex is an inherently complicated system with regards to both structure and function; hence improving prevention, detection, and treatment of glenohumeral pathologies has been difficult. Fortunately, shoulder biomechanics is an expanding and developing field that has and will continue to play an instrumental role in furthering the understanding of the anatomy and behavior of the shoulder complex and in enhancing the ability of medicine to detect and repair various pathologies. Advancements in the biomechanical analysis of the glenohumeral joint and in the ability of medicine to manage shoulder injuries are closely related, and both biomechanics and medicine have reaped considerable benefits from technological and scientific developments in medical imaging and surgical techniques that have enabled a new perspective of the glenohumeral joint with regards to both form and function.



<http://www.shoulderdoc.co.uk/images/uploaded/SLAP%20region.jpg>

Figure 1: SLAP Lesion Injury Region

The advent of shoulder arthroscopy as a medical tool led to the initial identification and description of glenoid labral tears in 1985, a glenohumeral

musculoskeletal pattern and abnormality that was previously undetectable<sup>1</sup>. The specific glenoid labral pathology, a superior labrum anterior-posterior tear was coined 'SLAP lesion' in 1990 (*Figure 1*)<sup>32</sup>. Although it has been more than two decades since SLAP lesions were defined and despite considerable advances in the understanding of glenohumeral biomechanics and glenoid labral pathologies, SLAP lesion detection remains difficult. Radiography and physical examination have proven useful for assessing a wide variety of orthopedic injury, but have shown limited potential with regards to SLAP lesion detection. Arthroscopy remains the 'gold standard' and the only definitive means of SLAP lesion diagnosis<sup>18, 22, 33 4, 8</sup>. An alternative to shoulder arthroscopy would be advantageous due to the invasiveness, financial cost, and patient risk associated with arthroscopy.

Recent developments in advanced imaging methods have drastically improved the diagnostic reliability of radiography, particularly in the detection of musculoskeletal patterns and injuries that previously, due to the low contrast of x-ray and computed tomography (CT), were impossible to identify with radiography. Although, magnetic resonance (MR) arthrography, specifically in high contrast, has shown some promise as a supplementary tool in SLAP lesion diagnosis, natural anatomic variants inherent in shoulder anatomy limit the reliability of radiographic diagnoses<sup>3, 4, 7, 21, 26, 35</sup>. Furthermore, problems with radiography follow those of arthroscopy; MR arthrography, for example, can be invasive, expensive, and dangerous, causing life threatening allergic reactions in some patients<sup>5</sup>. Therefore radiography is an imperfect means of SLAP lesion detection.

More than 20 provocative tests for the clinical evaluation of SLAP lesions are proposed in the literature. In most cases, the evaluation of these physical exams claim to have promising accuracy for the detection of SLAP lesions by their original authors<sup>2, 19, 24, 27, 36, 37</sup>. However, secondary studies fail to reproduce the initial findings, typically reporting much lower values for the sensitivity and specificity of the physical examination tests<sup>8, 11, 14, 22, 28, 29</sup>. The discrepancies between the findings of these studies most likely reflect two primary difficulties;

- 1) The clinical detection of SLAP lesions is hindered by the fact that SLAP lesions are rarely isolated; meaning they are frequently accompanied by other various glenohumeral pathologies which are potential sources for labral symptoms<sup>3</sup>.
- 2) Differences in study protocols and problems associated with the methods used to verify accuracy of the design of these tests make comparisons between studies and verification of SLAP lesion tests difficult<sup>8</sup>.

The bulk of the literature determines SLAP lesion test diagnostic accuracy utilizing a single verification method. Typically a patient with a suspected SLAP lesion performs the provocative tests of interest in a clinical setting before shoulder arthroscopy. The outcome of the SLAP lesion tests from the clinical evaluation is then verified with conclusive arthroscopic findings<sup>7-9, 11, 14, 22, 23, 29</sup>. The results of these comparative studies have significant quantitative discrepancies, but a fundamental qualitative conclusion recurs; no single SLAP lesion test has the sensitivity or specificity to independently determine the presence or absence of a SLAP lesion<sup>7-9, 28</sup>. Although previous studies attempt to assess the diagnostic accuracy of SLAP lesion tests, the analyses do little to

explain the reasons behind their apparent failure and rarely suggest or point to any means of improving the performance of the tests.

Although, clinically based evaluations of SLAP lesion tests account for the majority of studies to date, studies have also assessed test accuracy by attempting to validate the fundamental design behind various tests. Provocative SLAP lesion tests, by definition, function to provoke labral symptoms (primarily in the form of shoulder pain) as a positive diagnostic sign, by reenacting one of two injury mechanisms.




Figure 2: SLAP Lesion Injury Mechanism I

The first mechanism (*Figure 2*) elicits active tension in the biceps tendon and is typically associated with an acute traction trauma to the arm or elicited from repetitive overhead throwing injuries. The tensile load produced in the biceps tendon can pull and damage the superior labrum, the functional link between the insertion of the biceps tendon and the glenoid rim. The second injury mechanism produces passive compression of the humeral head and is often associated with a fall to outstretched arms (*Figure 3*).




Figure 3: SLAP Lesion Injury Mechanism II

The compressive load causes superior humeral head translation within the glenohumeral joint and can result in a collision between the humerus and labrum, potentially damaging the soft tissue of the labrum<sup>1</sup> The ability of proposed tests to reenact the injury mechanisms that they were designed to replicate has been examined from a few different perspectives including anatomic<sup>10, 17, 30</sup> and electromyographic<sup>10, 34</sup> methods. The results of these studies illuminate the importance of design validation during the development of clinical testing procedures.

The results of these studies do not clearly define the most accurate test for SLAP lesion diagnosis. Therefore the purpose of this study was to use electromyography (EMG) to biomechanically assess the ability of seven provocative tests to create active tension in the biceps tendon by activating the long head of the biceps brachii (LHBB). The SLAP lesion tests in this study were expected to successfully elicit LHBB activity because they were designed by their original author to reproduce a SLAP lesion injury mechanism in that manner; hence this study was a means of verifying the design of SLAP lesion tests that are meant to reproduce the first SLAP lesion injury mechanism. Furthermore, this study also examined six other glenohumeral joint stabilizing muscles,

the short head of the biceps brachii (SHBB), anterior deltoid (DELTA), pectoralis major (PECT), latissimus dorsi (LAT), infraspinatus (INFRA), and supraspinatus (SUPRA), with EMG to determine how effectively each test isolated the activation of the LHBB and to characterize the behavior of each of the glenohumeral joint stabilizing muscles.

Selectively activating the LHBB should reduce confounding implications of labral symptoms elicited from a source other than a SLAP lesion. Also, slight modifications were made to each of the original authors' depiction of the tests to allow the use of the Biodex System II Dynamometer to improve the uniformity between each subject's anatomical orientation and performance for each test. Additionally, the Biodex System aided the attempt to control and limit differences that may have resulted from inconsistencies in the magnitude of load applied for each test and the impact of the variability in subject's strength and their respective ability to resist the applied load.

The primary objective of this study was to evaluate seven SLAP lesion tests, chosen based on the findings of a pilot study, in subjects with no history of shoulder pathology, using EMG according to two variables, muscle activation and muscle selectivity, to characterize particular aspects of LHBB behavior. Additionally, differences in the LHBB behavior between male and female gender groups were examined. Furthermore, a brief supplementary analysis characterized the behavior of the six other joint stabilizing muscles in the same manner as for the LHBB.

The hypotheses of this study was that there would be no difference between the seven SLAP lesion tests in the LHBB behavior for either variable, no crucial behavior differences were anticipated between tests for the other six glenohumeral joint stabilizing

muscles, and furthermore differences between gender groups were also not anticipated.

The statistical hypotheses will be as follows:

$H_0 : \mu_1 = \mu_2 = \dots = \mu_7$  (All SLAP lesion tests elicit the same value in)

- LHBB activation
- LHBB selectivity
- Gender groups

$H_A : \mu_1 = \mu_2 = \dots = \mu_7$  (All SLAP lesion tests elicit a different value in)

- LHBB activation
- LHBB selectivity
- Gender groups

### **Operational Definitions**

This study involves two dependent variables that characterize a specific behavior of the individual muscles. The principal interest for SLAP lesion detection was the LHBB behavior, but regardless of the muscle analyzed, for each respective muscle the variables were quantified by the same method for each of the seven respective muscles.

- *LHBB activation*: the maximum LHBB activity elicited
  - as indicated by the peak LHBB amplitude of the normalized EMG signal recorded during each respective provocative tests, units of percent maximum contraction, range ideally between zero and 100%
- *LHBB Selectivity*: the ratio depicting the ability of a respective provocative test to selectively activate the LHBB

- as indicated by the ratio of the LHBB activation defined above, over the sum of all seven glenohumeral muscles respective activations, defined by the same method above but with respect to each muscle, the normalized maximum peak EMG signal amplitudes elicited during a respective provocative test, unitless due to the ratio of units of percent maximum contraction over units of percent maximum contraction, range between zero and one

### **Limitations**

The results of this study are limited to individuals who are represented by the sample population: healthy males and females who have no history of shoulder pathology. The findings in this study are only representative of individuals falling within the subject parameters noted above. It should be noted that patients with a suspected SLAP lesion may have considerable differences in muscle behavior than those who have had no history of shoulder pathology, like the cohort in this study.

### **Delimitations**


The results of this study are applicable to all physicians and clinicians using any of the seven tests because the focus is simply to verify the tests' design by assessing the ability of each test to reproduce a specific SLAP lesion injury mechanism. In elaboration, this study allows the verification of the test design, which is limited in other studies with subjects who have a shoulder injury as a successful test. The findings of such studies may not accurately represent the fundamental ability of the test to reproduce the injury mechanism.

## LITERATURE REVIEW

This chapter summarizes the findings of relevant journal articles that have been published regarding SLAP lesions to date. Topics include shoulder and labral anatomy, the history of SLAP lesions, and the literature containing evaluations of the seven provocative SLAP lesion tests of interest for the present study described in this thesis

### Shoulder Anatomy


The shoulder complex (*Figure 4*) is an intricate system containing four bones (clavicle, humerus, thorax, and scapula), three anatomical articulations (acromioclavicular, sternoclavicular, and glenohumeral), and one functional articulation (scapulothoracic), which is supported by ligamentous structures, soft tissues, and the musculature surrounding of the shoulder girdle.



<http://www.pt.ntu.edu.tw/hmchai/Kinesiology/KINupper/Shoulder.files/ShoulderStructure.htm>

Figure 4: Shoulder Complex - Bony Anatomy

The interaction of these components produces the most dynamic and mobile joint complex in the body. The stability of this uniquely mobile joint, and specifically the glenohumeral ball and socket joint (*Figure 5*), is maintained by a number of static and dynamic stabilizing structures. The articular surfaces of the proximal humerus and glenoid are mismatched with regards to size and orientation, which grants the joint extreme mobility, and essentially eliminates bony stability<sup>21</sup>.



<http://www.pt.ntu.edu.tw/hmchai/Kinesiology/KINupper/Shoulder.files/ShoulderStructure.htm>

Figure 5: Glenohumeral Joint Anatomy

Shoulder stability is maintained by a complex web of contributors including the static soft tissue structures of the joint itself and the musculature surrounding the shoulder girdle. A recent publication by Veeger and colleagues on shoulder biomechanics articulately noted that shoulder function is the ‘perfect compromise between stability and mobility’<sup>1</sup>. Clearly, the stability of the shoulder can be easily compromised due to the number of components and the complexity of their interactions. Glenoid labral pathologies can hinder the careful balance required by this unique biomechanical system. Glenoid labral musculature, surrounding connective tissues, and negative intra-articular

pressure are all proposed constituents involved in maintaining the stable position of the humeral head within the glenoid. The labrum itself plays a valuable role in joint stability. The glenoid labrum creates a suction effect on the humeral head and it increases the depth of the glenoid cavity by fifty percent. Hence, the presence of glenoid labral pathologies inherently affects stability of the glenohumeral joint.

### **SLAP Lesions**

In the last century, the development of medical imaging techniques has radically expanded the understanding of human anatomy. Radiography began with the discovery of the x-ray in the late nineteenth century, and its diagnostic value was quickly realized. Today the term radiography encompasses the range of imaging modalities not limited to computed tomography (CT) and magnetic resonance imaging (MRI). Both CT and MRI create three dimensional reconstructed images, which improve on the flat, two dimensional nature of the x-ray. MRI has far greater contrast than CT, which enables various soft tissues such as muscles and ligaments to be distinguished, greatly aiding the understanding of musculoskeletal anatomy. Magnetic resonance (MR) arthrography, where a contrast medium is injected into a joint or region of interest prior to MRI to further improve tissue differentiation, has been particularly useful for examining fine musculoskeletal pathologies including lesions of the shoulder<sup>4</sup>. Clearly, radiography has allowed human anatomy and physiology to be viewed in a new perspective, and specifically, advanced imaging methods have helped to illuminate the difficult form, function, and carefully balanced means of maintaining the stability of the shoulder complex.

The introduction of shoulder arthroscopy enabled the characterization of musculoskeletal pathologies that were previously unidentifiable with open surgical techniques, medical imaging modalities, or through the use of any other medical tools. In this manner, the complex musculoskeletal structure of the shoulder greatly benefited from the advent of shoulder arthroscopy. Specifically, in 1985 glenoid labral lesions were first described in throwing athletes after Andrews et al diagnosed 73 patients with the pathology after arthroscopic surgery. Andrews et al made several hypotheses based on observations during this early study, and they remain relevant today; 1) during throwing the biceps tendon undergoes large forces, 2) the most frequent location of glenoid labral tears is near the biceps tendon insertion at the anterior-posterior area of the glenoid (occurring in 83% of the glenoid labral lesion patients in the study by Andrews et al), and 3) the biceps tendon is likely the cause of glenoid labral lesion<sup>1</sup>.

In 1990, Snyder et al coined the term superior labrum anterior-posterior (SLAP) lesion to simply identify the labral lesion first described by Andrews et al<sup>32</sup>. Interestingly, although almost two decades has passed since this publication, the difficulty associated with SLAP lesions detection without arthroscopy, which was noted in the publication, has not changed considerably. Snyder et al examined more than 700 shoulders with arthroscopy and found 27 SLAP lesions. SLAP lesions were further categorized into four grades of severity ranging from Type I (where fraying and a general degenerative appearance of the superior aspect of the labrum is present) to Type IV (where 'bucket handle' tears are present and often are displaced into the joint with the lesion extending into the biceps tendon). Furthermore, importance of this study is indicated by the frequency with which it is referenced in the literature. Snyder et al was

the first to acknowledge the frequent occurrence of other shoulder pathologies with SLAP lesions, and the findings supported and improved upon concepts relating to SLAP lesion injury mechanisms. The injury mechanisms were related to either a tensile load on the biceps tendon or a compressive load received on the labrum itself. The injury occurred from either traction to the arm from a sudden traumatic increase in load, from a repetitive tensile load seen in overhead throwing athletes, or from a compressive load caused from a fall to outstretched arms.

An EMG study during baseball pitching noted that peak biceps brachii muscle activity occurred following ball release, during the deceleration phase of the arm while throwing<sup>15</sup>. These findings supported Andrews et al's biomechanical evaluation of throwing in the 1985 manuscript, examining the elbow and shoulder moments using three dimensional high speed cinematography and computer assisted analysis. Andrews determined that during the peak acceleration phase of throwing, the elbow extends from 80° to 30° in a 25-ms period, producing a peak moment of 600 inch-pounds prior to deceleration. The hypothesis is that the burst of biceps activity at the beginning of the deceleration phase indicates that the biceps play a role in decelerating the joint and that the large mechanical moment noted may be dampened and controlled by the LHBB, supporting the possibility that the deceleration phase of throwing may be a likely cause of SLAP lesions<sup>1</sup>.

The present findings and consensus in the literature continues to support the proposed SLAP lesion injury mechanisms suggested over twenty years ago in the first two SLAP lesion publications<sup>1,32</sup>, but contrarily there have been some unique case reports whose findings seem to question these mechanisms. Specifically, the role of

tension in the biceps tendon in SLAP lesions has been called into question. A case study by Keefe et al discussed the potential need to reevaluate the pathomechanics behind SLAP lesion mechanisms due a patient with a SLAP lesion (arthroscopic verification) that did not have a biceps tendon. The patient could not recall a traumatic traction event, a fall to outstretched arms, and clearly the injury could not have resulted due to tension in the biceps tendon during overhead throwing, though the subject was an active throwing athlete. Although a lack of the biceps tendon may not be common for the general population, this case study may imply that the role of the tendon in the deceleration phase of throwing may need to be reevaluated for the general population and for the tendon's part in SLAP lesions<sup>16</sup>.

### **Provocative Tests**

#### Active Compression Test

In 1988 O'Brien et al first proposed the Active Compression test. It was originally intended to assess acromioclavicular (AC) joint pathologies, but after anatomic validation using cadaver studies and following testing on subjects with suspected SLAP lesions, the authors claimed that the Active Compression test was useful for the clinical detection of SLAP lesions and various AC joint pathologies. The test was originally designed based on the description of a patient with a degenerative AC joint, who described the primary movements that reproduced his symptoms of pain. The author conducted a study of 318 patients with shoulder pain and reported promising findings. The results of the clinical tests were confirmed by either arthroscopic verification or radiography, and the findings alleged that the Active Compression test had 100% sensitivity, 99% specificity, a positive predictive value of 94.6%, and a negative

predictive value of 100%<sup>27</sup>. These results have not been reproduced in secondary studies, and one large potential source of error for the study is that medical imaging was used to verify the presence of SLAP lesion, and radiography is known to exhibit poor accuracy for SLAP lesion detection<sup>8</sup>. The Active Compression Test is one of the most evaluated SLAP lesion tests in the literature. Most studies indicate it performs with some promise, but in general high accuracy is not reported.

In response to questions associated with the initial study's findings and reliability of the accuracy for the Active Compression test, McFarland et al conducted a study with 426 patients all of which underwent arthroscopic confirmation and found 47% sensitivity, 55% specificity, a positive predictive value of 10%, and a negative predictive value of 91%<sup>22</sup>. This study clearly does not support previous findings, and again the discrepancy could be linked to the error associated with radiographic diagnoses and their use in the study.

Similar studies report findings which parallel the results of McFarland et al, including the 2001 study by Kim et al and 2006 study of Parentis et al. Both of the studies further categorized SLAP lesions into various subtypes, to assess potential improvements in clinical test performance when severity and type of SLAP lesion were taken into account. Unfortunately in both studies the accuracy of the Active Compression test was below 65 % regardless of the type of SLAP lesion<sup>19, 29</sup>.

Though much less in number, several other studies have attempted to evaluate the accuracy of the Active Compression SLAP lesion test utilizing different test design verification methods, and these analyses are of particular interest to the study in this thesis. A study in 2004 assessed the anatomical basis for the test, using MRI. The

findings suggested there was an anatomical basis for the test. The study suggests that the internal rotation of the arm, required for the palm-up position of the test, causes consistent physical contact between the superior labrum and lesser tuberosity, which likely would be a source of pain for subjects with a damaged labrum. Furthermore, this contact between articular surfaces is eliminated with internal rotation of the forearm as the palm-down position requires. These findings support the anatomic validation of the Active Compression Test because the test is considered positive for a SLAP lesion only when the patient has labral symptoms during the palm-up portion of the test that are not present during the palm-down portion of the test<sup>30</sup>.

In 2006 another study proposed that Type II SLAP lesions may be best detected with the Active Compression tests and EMG was used to determine which clinical tests in the study elicited the most promising muscle behavior. The study found that strong LHBB activity peak was elicited during the Active Compression test, indicating that the test may be a better diagnostic tool than other tests in that study<sup>34</sup>.

In 2008 a study also attempted to anatomically validate the Active Compression test using two methods. First, the objective was to quantify the active tension in the biceps tendon using EMG and twelve healthy subjects. Second, the objective was to quantify the passive tension in the tendon in five cadaver shoulders, using a custom designed load cell to determine strain on the biceps tendon. In contrast to Parentis et al, this study found that the anatomic basis of the Active Compression test was not valid<sup>10</sup>.

Although the Active Compression Test is examined frequently in the literature, the findings of these studies are limited, and this is a pattern that is repeated for the remaining six tests examined in this study. Majority of the studies assess this test, and all

SLAP lesion tests, by a single comparative method. The clinical findings of the test are compared with the concrete arthroscopic diagnoses, and such a comparative analysis is has considerable limitations. These comparative studies provide no indication as to why the test performed successfully or otherwise and furthermore, the tests provide no information or means of improving upon test performance. Studies that examine tests using other methods such as with EMG and cadaver specimens, improve upon the comparative analysis, in that they provide information that potentially explains reasons for their performance. In the case of the Active Compression test, biomechanical analyses were seen in two studies, using EMG and cadaver, but more studies in this manner would benefit SLAP lesion tests.

#### Speed's Test

In 1998, Speed's Test was introduced to assess a variety of shoulder pathologies, and this study is frequently used today in the clinical setting. Bennett et al assessed 46 shoulders in 45 patients with arthroscopic confirmation, and determined that Speed's had a promising sensitivity of 90%, but found the test performed poorly for other accuracy measures with 14% specificity, a 23% positive predictive value, and a 83% negative predictive value<sup>2</sup>. Another study countered these findings, eliciting low sensitivity results for Speed's tests at 32%, 75% specificity, a 50% positive predictive value, and a 58% negative predictive value. This study concluded that Speed's was moderately specific, but the test was unlikely to influence the pretest diagnosis held by the clinician. The authors reiterated the fallibility of clinical assessments, because depending on the setting and population, they argue that predictive values vary inherently<sup>14</sup>.

In 2007, another study examined the accuracy of Speed's to detect partial tears in the biceps tendon in 847 consecutive patients who underwent arthroscopy, and 40 of those had confirmed SLAP lesions. In this study, Speed's tests had a sensitivity of 50%, specificity of 67%, a positive predictive value of 8%, and negative predictive value of 96%. The frequent occurrence of other shoulder pathologies was attributed to the reason behind the poor behavior of Speed's and the anticipated unreliability of any clinical exam to detect partial tears of the biceps tendon. Of the 847 patients, 40 had partial bicep tendon tears, 34 had partial rotator cuff tears, three had anterior instability, two had impingement without rotator cuff tear, and 1 had degenerative arthritis. The study concluded that no single physical examination test can accurately predict the presence of a partial tear in the biceps tendon. The study also suggested that tests designed to produce tension in biceps tendon are not helpful in detecting partial tears of the bicep tendon<sup>9</sup>. Again the lack of valuable information that can be derived from these types of comparative studies must be reiterated, and Speed's has not been studied biomechanically or anatomically to date.

#### Pronated Load Test

The performance of the Pronated Load test has not been evaluated beyond the mention of promising sensitivity by Wilk et al in 2005. The test was designed to simulate the injury mechanism and peel back behavior seen during stimulation. The Pronated Load test is meant to have the promising behavior of the Pain Provocation Test which causes passive external rotation of the forearm, coupled with a position that enables large activity from the LHBB during contraction<sup>36</sup>. No biomechanical studies are presently available on the Pronated Load test.

### Bicep Load I Test

In 1999, a new SLAP lesion test was proposed; the Bicep Load I Test. This provocative test was designed to detect SLAP lesions in subjects who have recurrent anterior shoulder dislocation, typically found in SLAP lesion Type II subjects. The original and only study evaluating this test was a cohort study, and evaluated 75 patients who all underwent arthroscopic surgery. The Bicep Load I test indicated that 12 subjects had SLAP lesions, and 10 of these were arthroscopically confirmed as Type II SLAP lesions. The resulting test sensitivity was 90.9%, specificity was 96.9%, positive predictive value was 83.0% and negative predictive value was 98.0%<sup>20</sup>. Although, the original findings are promising, the test is designed for SLAP lesion detection only in shoulders with recurrent dislocation and may not be as reliable for those patients without the additional shoulder pathology. Furthermore, no biomechanical or anatomic studies have evaluated this test.

### Bicep Load II Test

In 2001, another SLAP lesion test was proposed, Bicep Load II test, as a complement to Bicep I. The Bicep Load II test was designed with the intent to detect isolated SLAP lesions. 127 subjects were evaluated in the study, 38 were positive for a SLAP lesion according to the Bicep Load II test, and 35 were confirmed to have SLAP lesions following arthroscopy. Again, promising accuracy was reported with 89.7% sensitivity, 96.6% specificity, 92.1% positive predictive value, and 95.5% negative predictive value<sup>19</sup>. These findings may be limited to isolated SLAP lesions, which is inherently uncommon.

In 2008, first study evaluating the ability of a combination of more than one provocative tests to detect Type II SLAP lesions was published. Several tests, including the Bicep Load II test, were included in this study, and interestingly the Bicep Load II test was categorized as a high performing test with respect to specificity. This study found that the combined findings from two relatively sensitive and one relatively specific test improved SLAP lesion detection accuracy dramatically, such that sensitivity was a minimum of 70% when one of the three tests were positive and specificity was a minimum of 90% when all three SLAP lesion tests were positive. Furthermore, in this study, the author explicitly stated that no single SLAP lesion test would have the capability, with regards to simultaneous strength in sensitivity and specificity, to be individually able to detect a Type II SLAP lesion<sup>28</sup>. Bicep Load II is another SLAP lesion test that has not been evaluated by means other than comparative assessment.

#### Resisted Supination External Rotation Test

In 2005, the Resisted Supination External Rotation was developed to mimic the peel-back mechanism associated with SLAP lesions. The study examined 40 athletes, of which 29 had SLAP lesions verified by arthroscopy. The results from the Resisted Supination External Rotation test were compared to those of the Crank test and the Active Compression test. Meyers et al claimed the Resisted Supination External Rotation test has better performance than both of the others, with a sensitivity of 82.8%, a specificity of 81.8%, a positive predictive value of 92.3%, and a negative predictive value of 64.3%<sup>24</sup>. Further evaluation of this test is needed, as the only study evaluating the test is this original study, and no biomechanical studies have been published to date.

### Supination Sign Test

The Supination Sign test another provocative test designed to elicit labral symptoms as a means of SLAP lesion detection that is anatomically based. The Supination Sign Test was shown to have specificity and low sensitivity across four studies including Nakagawa et al, Guanche et al, Holtby et al, and Parentis et al resulting in comparable finding for sensitivity (14%, 12%, 43%, and 13% respectively for each study) and similarly for specificity (98%, 96%, 79%, and 93%). Although in general the Supination Sign test has a high specificity, high specificity is likely not a good method for stand alone evaluation of the presence of a SLAP lesion<sup>11 14 25 29</sup>. Once again, no biomechanical assessment has been done.

In summary, no study has biomechanically assessed the accuracy and performance of these tests. Although many studies have attempted to determine test accuracy by comparative analysis and some studies have examined a single test anatomically or with EMG, further biomechanical assessment is necessary to properly evaluate the ability of these tests to aid in the detection of SLAP lesions in the clinical setting. A biomechanical evaluation of these tests, will not only help to verify the design of these test and provide an alternative method to quantify test accuracy, but biomechanical assessment could also provide valuable information as to how these tests may be improved.

## METHODS

This chapter addresses the methodology and procedures that were used to acquire the data necessary to fulfill the purpose of this study. Topics that will be addressed in this chapter include the experimental protocol for this study, the original descriptions of each provocative SLAP lesion test and the modifications used in this study, the methods used to filter and analyze the EMG data, including the definitions and mathematical equations for muscle activation and muscle selectivity, and the statistical methods used to determine the significance of the data.

### **Experimental Protocol**

#### Subject and IRB approval

A cohort of 21 healthy volunteers comprised of 11 females ( $24.7 \pm 6.7$  years,  $168.4 \pm 5.3$ cm,  $66.9 \pm 9.1$ kg) and 10 males ( $29.4 \pm 10.6$  years,  $178.1 \pm 6.6$  cm,  $80.0 \pm 6.4$ kg) with right arm dominance and no history of shoulder pathology were recruited for subjects in this study. Subjects recruited were either college students at Boise State University or medical health professionals from the local area. All procedures were approved by the Institutional Review Board at Boise State University, and all participants read and signed a statement of informed consent prior to the start of testing.

#### Electromyography Apparatus and Subject Preparation

EMG was used to record muscle activity for seven muscles surrounding the dominant arm's glenohumeral joint including the long head of the biceps brachii (LHBB), short head of biceps brachii (SHBB), anterior deltoid (DELTA), pectoralis major (PECT),

latissimus dorsi (LAT), infraspinatus (INFRA), and supraspinatus (SUPRA). Each subject was instrumented with a single 44-gauge fine-wire indwelling electrode and six surface bipolar silver-silver chloride EMG electrodes (Noraxon, USA Inc, Scottsdale, AZ). The surface electrodes were positioned over the muscle belly and parallel with the orientation of the muscle fibers, as seen below where a) LHBB and SHBB, b) DELT, c) PECT, d) LAT, e) INFRA, and f) SUPRA (*Figure 6*)<sup>6</sup>.




Figure 6: Electrode Placement for EMG (a – f)

An additional surface electrode was placed on the acromion process of the non-dominant shoulder to serve as a reference. Due to the location of the SUPRA deep to the trapezius, the indwelling electrode was necessary to acquire SUPRA activity. Using sterile techniques a certified medical technician placed the fine-wire indwelling electrode using a 27-gauge sterile needle. EMG data were recorded using the Vicon Nexus Software (Vicon, Los Angeles, CA) at 1250 Hz using a Noraxon Telemetry 900 EMG system (Noraxon USA, Inc, Scottsdale, AZ).

#### Subject Protocol

Using established EMG protocols, each subject was asked to perform Maximum Voluntary Isometric contractions (MVICs) for each of the seven muscles in random order on a Biodex System II Dynamometer (Biodex Medical Systems, Shirley, NY). Modifications were made to the MVIC recommendations of Cram, Hintermeister, and Rowlands<sup>6, 12, 31</sup> to accommodate for the use of the Biodex System (*Table 1*).

<b>Muscle</b>	<b>Joint Position</b>	<b>Resisted Maneuver</b>
<b>DELT</b>	Arm at side	Shoulder flexion
<b>LHBB / SHBB</b>	Elbow flexed 90°, shoulder flexed 90°	Elbow flexion
<b>INFRA</b>	Arm abducted 45°, elbow flexed 90°	External Rotation
<b>LAT</b>	Shoulder flexed 90°, arm internally rotated	Shoulder extension
<b>PECT</b>	Arm abducted 90°, forearm supinated	Horizontal adduction
<b>SUPRA</b>	Arm abducted 90°, forward flexed 30°, and internally rotated	Maintain against resistance

Table 1. MVIC Joint Positions and Resisted Maneuvers

For each trial six seconds of data were recorded to ensure that the entire burst of muscle activity was captured during each MVIC. The subjects performed three trials for each MVIC and were asked to maximally contract for a count of three seconds. Subjects rested for thirty seconds between MVIC trials to avoid fatigue effects. For each MVIC, the peak amplitude of the EMG signal among the three trials was used to normalize the provocative test data to a percentage of effort.

Similarly seven provocative tests were performed in random order based on the descriptions of the original authors but with modifications to accommodate for use of the Biodex System. These tests were chosen based on the findings of a preliminary pilot study that evaluated clinical tests from relevant literature that were designed to reenact SLAP lesion injury mechanisms. Again, each subject performed three trials for a three second count for each test, six seconds of data were recorded for each trial, and the subjects rested for thirty seconds between trials to avoid fatigue affects.

Once all MVIC and SLAP lesion test trials had been completed the surface electrodes were removed from the subject, and a trained medical technician, using sterile techniques, removed the indwelling electrode from the SUPRA by applying gentle and steady traction to the leads. A sterile bandage and pressure were applied to the location where the indwelling electrode was removed. Each subject was advised to seek medical attention if an infection appeared to develop at the site, although infection was not anticipated.

### **Test Descriptions and Modifications**

Each of the seven SLAP lesion tests performed in this study were provocative tests that were designed by their respective original authors to reproduce one SLAP

lesion injury mechanism, by activating the LHBB to induce tension in the biceps tendon. For the purposes of this study, modifications were possible and made for six of the seven SLAP lesion tests utilizing the Biodex System II Dynamometer to improve standardization between subjects and hopefully to reduce the potential for differences in muscle behavior or test performance due to variable subject and clinician strengths that could alter the intended test position or function. Tests requiring static resistance against an applied load maintained the static subject position through the stationary preset up of the Biodex System in isometric mode. Tests requiring dynamic resistance to an applied load were controlled by the Biodex System allowing a motion with a constant velocity regardless of the force applied by the subject through the isokinetic mode of the Biodex System. The original description of each test and the modifications employed in this study are noted below.

#### Active Compression Test (ACPU and ACPD)

The Active Compression Test has two positions, palm-down (ACPD) and palm-up (ACPU), which vary only by internal or external rotation of the arm. The patient is standing with the elbow in full extension, the shoulder is flexed to  $90^\circ$ , and adducted  $10-15^\circ$  medial to the sagittal plane. For ACPD (*Figure 7*) the arm is maximally internally rotated such that the thumb points down. The patient is asked to resist a uniform downward load applied to their arm by the clinician. For ACPU (*Figure 8*) the initial patient positioning is unchanged except the arm is externally rotated such that the palm faces up. Again, the patient is asked to resist a uniform downward load applied by the clinician<sup>27</sup>.



Figure 7: ACPU



Figure 8: ACPD

For the purpose of analysis and due to the nature of the subject population, a control group having no history of shoulder pathology, this study treated ACPU and ACPD as two independent tests. Both tests were modified such that the subject was seated in the Biodex System. The orientation of the subject's arm remained true to original description of the two test positions by O'Brien et al, but the subject was

asked to resist the stationary set-up of the arm of the Biodex System by attempting to lift his/her arm superiorly for both ACPD and ACPU.

### Speed's Test

According to the original description of Speed's test (Speed), the patient is standing and resists a downward force applied to the upper extremity with the elbow extended, forearm supinated, and arm elevated to  $90^{\circ}$ <sup>2</sup>. In this study the orientation of the subject's arm remained similar to the original definition, but the test was modified into a dynamic movement controlled by the Biodex System. The subject's arm started hanging beside and parallel to the body with the palm facing up (*figure 9*), and then the subject was asked to raise the arm (flex the shoulder) with as much force as possible to  $90^{\circ}$ . Regardless of the force exerted by the subject, motion was restricted to a constant velocity by the Biodex System in the isokinetic setting of  $60^{\circ}$  per second.



Figure 9: Speeds Starting Position

### Pronated Load Test

In the original description, the patient is in the seated position with elbow flexed to  $90^\circ$ , the arm abducted to  $90^\circ$ , maximally externally rotated, and the forearm is fully pronated. The patient is then asked to perform an isometric contraction of the biceps (a 'curl' with the forearm pronated)<sup>36</sup>. The Pronated Load test (ProLoad) was negligibly modified for this study. The subject sat in the original orientation in the Biodex System, the subject's arm was supported just proximal to the elbow. The subject was asked to perform an isometric bicep contraction (pronated curl) which was resisted by the static set up of the Biodex System.

### Bicep Load I Test



Figure 10: Bicep Load I

According to the original author's definition of the Bicep Load I test (Bicep I), the patient is in the supine position when an anterior apprehension test is performed starting with the arm abducted  $90^\circ$  with the forearm fully supinated<sup>20</sup>. Bicep I was modified such that the patient was seated in the Biodex System, in the same position

as ProLoad, except the forearm was fully supinated (*Figure 10*). The subject was asked to perform a bicep contraction (a traditional ‘curl’), which was resisted by the static set up by the Biodex System.

### Bicep Load II Test

The patient is supine with the arm abducted to 120° degrees, the elbow flexed to 90° degrees, and the forearm fully supinated.



Figure 11: Bicep Load II

The patient is then asked to flex the elbow against the resistance of the clinician<sup>19</sup>. For this study the modification for Bicep Load II test (Bicep II) paralleled those made to Bicep I except the arm was abducted to 120° degrees instead of 90° (*Figure 11*).

### Resisted Supination External Rotation Test

The original authors describe putting the patient in the supine position with scapula near the edge of an evaluation table; the patient’s arm is supported by the physician at the wrist, with the arm abducted to 90° and the elbow flexed between 65° and 70° degrees. The clinician then externally rotates the arm while the patient is

asked to supinate the forearm<sup>24</sup>. The Resisted Supination External Rotation test was essentially unchanged for this study (*Figure 12*), and the movement was not controlled by the Biodex System because the position and motion could not be recreated with the Biodex System for all subjects. One certified athletic trainer performed RSER with each subject in the study in the supine position on the Biodex System.




Figure 12: RSER Starting Position

### Supination Sign Test

As originally defined, the Supination Sign test (Yergason), in the seated position with the elbow flexed to 90° and forearm fully pronated, the patient is asked to attempt to supinate the forearm while the physician resists the attempted motion while holding the wrist<sup>37</sup>. Yergason was scarcely modified in this study; as the patient maintained the defined orientation but the forearm was fastened to the static arm of the Biodex System (*Figure 13*). The subject was asked to attempt to supinate the forearm against the static setup of the Biodex System.



Figure 13: Yergason's

### **Electromyography Analysis**

The raw EMG signals were filtered, normalized, and then analyzed to characterize muscle behavior using custom MATLAB software. The raw EMG signals were processed a traditional EMG filtering technique that is frequently employed with EMG and noted in the literature. After processing the EMG signals, numerical values were calculated for the LHBB activation and LHBB selectivity as a means of characterizing LHBB behavior. Supplementary calculations were made for the muscle activations and muscle selectivities of the remaining six glenohumeral muscles in this study.

#### **Initial EMG Processing**

During all testing, EMG data from each muscle were acquired at 1250 Hz, and the raw data was band-pass filtered from 16 to 500 Hz by the data collection unit internally prior to transmission of the data to the wireless receiver for further processing. Next, custom MATLAB software was used to further process, normalize, and analyze the EMG signals.

### EMG Filtering Technique

The majority of EMG signals that were collected, the LHBB, SHBB, DELT, PECT, and SUPRA, were filtered by a traditional smoothing technique and band-pass filtered from 20 to 500 Hz, and then the signals were rectified and smoothed using a root mean square algorithm in combination with a 20-ms forward moving window average.

### Normalizing Provocative Test Data with MVIC Maximums

The provocative test EMG signals were normalized to a percentage of effort based on the peak EMG amplitude elicited during each muscle's respective Maximum Voluntary Isometric Contraction (MVIC). Ideally the normalized and filtered EMG signals for each provocative test would have a muscle activation range of zero to 100% MVIC. Muscle activation and muscle selectivity were calculated for each muscle during each test.

### Muscle Activation

Muscle activation was used to determine how effective each SLAP lesion test was at causing the individual muscles to activate and was defined as the peak muscle activities elicited during the three normalized trials.

### Muscle Selectivity

The ability of each provocative test to isolate the LHBB is important for diagnosing SLAP lesions due to the common association of SLAP lesions with other shoulder pathologies. Therefore in this study, a ratio depicting the ability of each test to selectively activate each muscle was calculated. The muscle selectivity for each test was defined as the ratio of the peak activation of the muscle of interest over the sum of peak activations for all seven muscles examined in the study. For example, a selectivity ratio

of 1.0 for any muscle,  $N$ , would indicate that muscle  $N$  was the only active muscle contributing to the EMG signal while the other six muscles remained inactive. For each test the selectivity for muscle  $N$  was calculated using the following equation.

$$N\_SelectivityRatio = \frac{A_N}{\sum (A_{LHBB} + A_{SHBB} + A_{DELT} + A_{PECT} + A_{LAT} + A_{INFRA} + A_{SUPRA})}$$

- where  $N$  is the muscle of interest (LHBB, SHBB, DELT, PECT, LAT, INFRA, or SUPRA)
- where  $N\_SelectivityRatio$  is the selectivity ratio of muscle  $N$
- where  $A_N$  is the peak muscle activation for muscle  $N$

### **Statistical Analysis**

Statistical analysis was conducted using SPSS Statistics Software 17.0 for Windows to determine significance of the data, specifically, differences in maximum muscle activations and muscle selectivities for each test, between tests, and between male and female groups. A repeated measures analysis of variance (ANOVA) test was performed to identify significant differences between provocative tests for each individual muscle. A pair-wise T-test post-hoc analysis was performed to compare results between each test using a p-value sliding scale Bonferroni adjustment<sup>13</sup>. Likewise, a paired-sample T-test was used to examine potential differences in muscle activation ( $p = 0.05$ ) between male and female groups.

## REFERENCES

1. Andrews, J., W. Carson, Jr, and W. McLeod, *Glenoid labrum tears related to the long head of the biceps*. Am J Sports Med, 1985. **13**(5): p. 337-341.
2. Bennett, W.F., *Specificity of the Speed's Test: Arthroscopic Technique for Evaluating the Biceps Tendon at the Level of the Bicipital Groove*. The Journal of Arthroscopic and Related Surgery, 1998. **14**(8): p. 789-796.
3. Chandnani, V., et al., *Glenoid labral tears: prospective evaluation with MRI imaging, MR arthrography, and CT arthrography*. Am. J. Roentgenol., 1993. **161**(6): p. 1229-1235.
4. Chang, D., et al., *SLAP lesions: Anatomy, clinical presentation, MR imaging diagnosis and characterization*. European Journal of Radiology, 2008. **68**(1): p. 72-87.
5. Chundru, U., *Magnetic Resonance Arthrography*. Musculoskeletal, 2009.
6. Cram, J.R. and G.S. Kasman, *Introduction to surface electromyography*. 1998, Gaithersburg, MD: Aspen Publishers Inc.

7. DaSilva, J.J., A.G. McNickle, and B.J. Cole, *Arthroscopic Superior Labrum Anterior to Posterior Repair*. Operative Techniques in Orthopaedics, 2008. **18**(1): p. 53-61.
8. Dessaur, W. and M. Magarey, *Diagnostic accuracy of clinical tests for superior labral anterior posterior lesions: a systematic review*. Journal of Orthopaedic and Sports Physical Therapy, 2008. **38**(6): p. 341-352.
9. Gill, H.S., et al., *Physical Examination for Partial Tears of the Biceps Tendon*. Am J Sports Med, 2007. **35**(8): p. 1334-1340.
10. Green, R.A., et al., *An evaluation of the anatomic basis of the O'Brien active compression test for superior labral anterior and posterior (SLAP) lesions*. Journal of Shoulder and Elbow Surgery, 2008. **17**(1): p. 165-171.
11. Guanche, C.A. and D.C. Jones, *Clinical Testing for Tears of the Glenoid Labrum*. The Journal of Arthroscopic and Related Surgery, 2003. **19**(5): p. 517-523.
12. Hintermeister, R.A., et al., *Electromyographic Activity and Applied Load During Shoulder Rehabilitation Exercises Using Elastic Resistance*. The American Journal of Sports Medicine, 1998. **26**(2): p. 210-220.
13. Holm, S., *A simple sequentially rejective multiple test procedure*. Scandinavian Journal of Statistics, 1979. **6**: p. 65-70.
14. Holtby, R. and H. Razmjou, *Accuracy of the Speed's and Yergason's Tests in Detecting Biceps Pathology and SLAP Lesions: Comparison With Arthroscopic Findings*. The Journal of Arthroscopic and Related Surgery, 2004. **20**(3): p. 231-236.

15. Jobe, F. and D. Moynes, *Delineation of diagnostic criteria and a rehabilitation program for rotator cuff injuries*. Am J Sports Med, 1982. **10**: p. 336-339.
16. Keefe, D.T., Lowe, WR, *Symptomatic Superior Labral Anterior Posterior Lesion With Absence of the Long Head of the Biceps Tendon*. The American Journal of Sports Medicine, 2005. **33**(11): p. 1746-1750.
17. Keren J. Shanley, R.A.G., Nicholas F. Taylor,, *An evaluation of the anatomical basis of the Biceps Load tests I and II for superior labral anterior and posterior (SLAP) lesions*. Clinical Anatomy, 2008. **21**(7): p. 647-651.
18. Kibler, W.B., *Specificity and Sensitivity of the Anterior Slide Test in Throwing Athletes With Superior Glenoid Labral Tears*. The Journal of Arthroscopic and Related Surgery, 1995. **11**(3): p. 296-300.
19. Kim, S.-H., et al., *Biceps Load Test II: A Clinical Test for SLAP Lesions of the Shoulder*. The Journal of Arthroscopic and Related Surgery, 2001. **17**(2): p. 160-164.
20. Kim, S.-H., K.-I. Ha, and K.-Y. Han, *Biceps Load Test: A Clinical Test for Superior Labrum Anterior and Posterior Lesions in Shoulders With Recurrent Anterior Dislocations*. The American Journal of Sports Medicine, 1999. **27**(3): p. 300-303.
21. Lugo, R., P. Kung, and C.B. Ma, *Shoulder biomechanics*. Shoulder Imaging European Journal of Radiology, 2008. **68**(1): p. 16-24.
22. McFarland, E.G., T.K. Kim, and R.M. Savino, *Clinical Assessment of Three Common Tests for Superior Labral Anterior-Posterior Lesions*. The American Journal of Sports Medicine, 2002. **30**(6): p. 810-815.

23. Munro, W. and R. Healy, *The validity and accuracy of clinical tests used to detect labral pathology of the shoulder* “ A systematic review. *Manual Therapy*. **In Press, Corrected Proof**.
24. Myers, T.H., J.R. Zemanovic, and J.R. Andrews, *The Resisted Supination External Rotation Test: A New Test for the Diagnosis of Superior Labral Anterior Posterior Lesions*. *The American Journal of Sports Medicine*, 2005. **33**(9): p. 1315-1320.
25. Nakagawa S, Y.M., Hayashida K, Obata M, Fukushima S, Miyazaki Y., *Forced shoulder abduction and elbow flexion test: a new simple clinical test to detect superior labral injury in the throwing shoulder*. *Arthroscopy*, 2005. **21**: p. 1290-1295.
26. Nam, E.K. and S.J. Snyder, *The diagnosis and treatment of superior labrum, anterior and posterior (SLAP) lesions*. *The American Journal of Sports Medicine*, 2003. **31**(5): p. 798-810.
27. O'Brien, S.J., et al., *The Active Compression Test: A New and Effective Test for Diagnosing Labral Tears and Acromioclavicular Joint Abnormality*. *The American Journal of Sports Medicine*, 1998. **26**(5): p. 610-613.
28. Oh, J.H., et al., *The Evaluation of Various Physical Examinations for the Diagnosis of Type II Superior Labrum Anterior and Posterior Lesion*. *Am J Sports Med*, 2008. **36**(2): p. 353-359.
29. Parentis, M.A., et al., *An Evaluation of the Provocative Tests for Superior Labral Anterior Posterior Tears*. *Archives of Physical Medicine and Rehabilitation*, 2005. **34**(2): p. 1-4.

30. Parentis, M.A., et al., *An Anatomic Evaluation of the Active Compression Test*. Journal of Shoulder and Elbow Surgery, 2004. **13**(4): p. 410-416.
31. Rowlands, L., et al., *Kinesiology of the Empty Can Test*. Am J Phys Med Rehabil, 1995. **74**(4): p. 302-304.
32. Snyder, S.J., et al., *SLAP lesions of the shoulder*. Arthroscopy: The Journal of Arthroscopic & Related Surgery, 1990. **6**(4): p. 274-279.
33. Stetson, W.B. and K. Templin, *The Crank Test, the O'Brien Test, and Routine Magnetic Resonance Imaging Scans in the Diagnosis of Labral Tears*. The American Journal of Sports Medicine, 2002. **30**(6): p. 806-809.
34. Swaringen, J.C., et al., *Electromyographic analysis of physical examination tests for type II superior labrum anterior-posterior lesions*. Journal of Shoulder and Elbow Surgery, 2006. **15**(5): p. 576-579.
35. Veeger, H.E.J. and F.C.T. van der Helm, *Shoulder function: The perfect compromise between mobility and stability*. Journal of Biomechanics, 2007. **40**(10): p. 2119-2129.
36. Wilk, K.E., et al., *Current Concepts in the Recognition and Treatment of Superior Labral (SLAP) Lesions*. Journal of Orthopaedic and Sports Physical Therapy, 2005. **35**: p. 273-291.
37. Yergason, R.M., *Supination Sign*. 1995.

APPENDIX

**Manuscript for Submission to the Journal of Shoulder and Elbow Surgery**

1           Abstracts for this study have been submitted and accepted for presentation at two  
2 annual conferences. First, the study will be presented as a podium at the 2009 Northwest  
3 Biomechanics Symposium (NWBS) at Washington State University, June 5<sup>th</sup> and 6<sup>th</sup> in  
4 Pullman, WA. Also, the study will be presented at the 2009 Annual Meeting of the  
5 American Society of Biomechanics (ASB) at Pennsylvania State University, August 26<sup>th</sup>  
6 through 29<sup>th</sup> in State College, PA. This manuscript has not been submitted for journal  
7 publication; furthermore no portion of the data, methods, results, or findings from this  
8 work have been previously submitted, published, or printed with the exception of the  
9 abstracts for the conference submissions noted above.

10           This manuscript has been read and approved by all authors, and each author  
11 believes the manuscript to be honest work. Dr. Michelle B. Sabick should be noted as  
12 the corresponding author for future inquiry with regards to this manuscript; her contact  
13 information is as follows:

14

15   **Corresponding Author:** Dr. Michelle B. Sabick, PhD

16   **Address:** Boise State University, Department of Mechanical and Biomedical Engineering,  
17 ET 204, 1910 University Dr., Boise, ID 83725-2075

18   **Office Phone Number:** 208.426.5653

19   **Fax Number:** 208.426.4800

20   **E-mail Address:** msabick@boisestate.edu

21

22 Vanessa J.C. Wood, MS, \_\_\_\_\_

23         This author, their immediate family, and any research foundation with which they  
24 are affiliated did not receive any financial payments or other benefits from any  
25 commercial entity related to the subject of this article.

26

27 Michelle B. Sabick, PhD, \_\_\_\_\_

28         This author, their immediate family, and any research foundation with which they  
29 are affiliated did not receive any financial payments or other benefits from any  
30 commercial entity related to the subject of this article.

31

32 Ron P. Pfeiffer, EdD, \_\_\_\_\_

33         This author, their immediate family, and any research foundation with which they  
34 are affiliated did not receive any financial payments or other benefits from any  
35 commercial entity related to the subject of this article.

36

37 Seth M. Kuhlman, MS, \_\_\_\_\_

38         This author, their immediate family, and any research foundation with which they  
39 are affiliated did not receive any financial payments or other benefits from any  
40 commercial entity related to the subject of this article.

41

42 Jason H. Christensen, \_\_\_\_\_

43           This author, their immediate family, and any research foundation with which they  
44 are affiliated did not receive any financial payments or other benefits from any  
45 commercial entity related to the subject of this article.

46

47 Mike J. Curtin, MD, \_\_\_\_\_

48           This author, their immediate family, and any research foundation with which they  
49 are affiliated did not receive any financial payments or other benefits from any  
50 commercial entity related to the subject of this article.

51

52           This project was supported by grant number 1-R15 AR054637 through the National  
53 Institute of Arthritis and Musculoskeletal and Skin Diseases. Grant funding was used to  
54 cover the overhead cost of the study, and to support a graduate research student who  
55 worked primarily with this study who was involved in data collection, data analysis, and  
56 manuscript preparation while under the guidance of the Co-Principal Investigators, Drs.  
57 Michelle Sabick and Ron Pfeiffer.

58           The study was approved by the Institutional Review Board at Boise State University,  
59 approval number 130.05.018, and each subject signed a form of informed consent prior to  
60 their participation in the study.

61

62 *Title Page*

63 **GLENOHUMERAL MUSCLE ACTIVATION**  
64 **DURING PROVOCATIVE TESTS DESIGNED TO DIAGNOSE SUPERIOR**  
65 **LABRUM ANTERIOR-POSTERIOR LESIONS**

66 <sup>1,2</sup>Vanessa J.C. Wood, MS, <sup>1,2</sup>Michelle B. Sabick, PhD, <sup>1,3</sup>Ron P. Pfeiffer, EdD, ATC,

67 LAT<sup>1,2</sup>, Seth M. Kuhlman, MS, <sup>1,2</sup>Jason H. Christensen, <sup>4</sup>Mike J. Curtin, MD

68 <sup>1</sup>Center for Orthopaedic & Biomechanics Research, Boise State University

69 <sup>2</sup>Department of Mechanical and Biomedical Engineering, Boise State University

70 <sup>3</sup>Department Kinesiology, Boise State University

71 <sup>4</sup>Intermountain Orthopaedics, Boise, Idaho

72

73 **Corresponding Author:** Dr. Michelle B. Sabick, PhD

74 **Address:** Boise State University, Department of Mechanical and Biomedical Engineering,

75 ET 204, 1910 University Dr., Boise, ID 83725-2075

76 **Office Phone Number:** 208.426.5653

77 **Fax Number:** 208.426.4800

78 **E-mail Address:** msabick@boisestate.edu

79

80 **IRB Approval:** Institutional Review Board at Boise State University (130.05.018)

81 **Grant Support:** National Institute of Arthritis and Musculoskeletal and Skin Diseases

82 (1-R15 AR054637)

83 **Disclaimer:** None

84 **Abstract**85 **BACKGROUND**

86 Despite considerable medical advances, arthroscopy remains the only definitive means of  
87 Superior Labrum Anterior-Posterior (SLAP) lesion diagnosis. Natural shoulder anatomic  
88 variants limit the reliability of radiographic findings and clinical evaluations are not  
89 consistent. Accurate clinical diagnostic techniques would be advantageous due to the  
90 invasiveness, patient risk, and financial cost associated with arthroscopy. The purpose of  
91 this study was to examine the behavior of the joint stabilizing muscles in promising  
92 provocative tests for SLAP lesions. Electromyography was used to characterize the  
93 muscle behavior, with particular interest in the long head biceps brachii, as activation of  
94 the long head and subsequent tension in the biceps tendon should elicit labral symptoms  
95 in SLAP lesion patients.

96

97 **METHODS**

98 Volunteers (N=21) without a history of shoulder pathology was recruited for this study.  
99 The tests analyzed were Active Compression, Speed's, Pronated Load, Biceps I, Biceps II,  
100 Resisted Supination External Rotation, and Supination Sign. Tests were performed on a  
101 dynamometer to improve reproducibility. Muscle activity was recorded for the long and  
102 short heads of the biceps brachii, anterior deltoid, pectoralis major, latissimus dorsi,  
103 infraspinatus, and supraspinatus. Muscle behavior for each test was characterized by peak  
104 activation and selectivity.

105

106 **RESULTS**

107 Speed's, Active Compression Palm-Up, Bicep I and Bicep II, produced higher long head  
108 activations. Resisted Supination External Rotation, Bicep I, Bicep II, and Yergason's,  
109 produced higher long head selectivities.

110

111 **CONCLUSION**

112 Bicep I, and Bicep II elicited promising long head behavior (high activation and  
113 selectivity). Speed's and ACPU elicited large long head activity, and Resisted Supination  
114 and Yergason's elicited selective long head activity. These top performing tests utilize a  
115 unique range of test variables that may prove valuable for optimal SLAP test design and  
116 performance.

117

118 **LEVEL OF EVIDENCE:** Diagnostic Study Level I

119

120 **KEY WORDS:** SLAP, Superior Labrum Anterior Posterior Lesion, provocative test,  
121 long head biceps brachii, diagnoses

122

123 ***Introduction.***

124           The advent of shoulder arthroscopy as a medical tool led to the description of  
125 glenoid labral tears in 1985<sup>1</sup>, and superior labrum anterior-posterior tears were coined  
126 ‘SLAP lesions’ in 1990<sup>30</sup>. Although it has been almost two decades since SLAP lesions  
127 were defined, diagnosis remains difficult<sup>17, 21, 31</sup> despite considerable advances in the  
128 understanding of glenohumeral biomechanics and glenoid labral pathologies. In spite of  
129 these advances, arthroscopy remains the only definitive means of SLAP lesion detection<sup>4,</sup>  
130<sup>8</sup>. Accurate clinical diagnostic techniques, as an alternative to shoulder arthroscopy,  
131 would be advantageous due to the invasiveness, financial cost, and patient risk associated  
132 with arthroscopy.

133           Radiography and physical examination have proven useful for assessing a wide  
134 variety of orthopedic injury, but have shown limited potential with regards to SLAP  
135 lesion detection. Though radiography, particularly MR arthrography in high contrast, has  
136 shown some promise as a supplementary tool in SLAP lesion diagnosis, natural anatomic  
137 variants limit the reliability of all radiographic diagnoses. Furthermore, parallel to  
138 arthroscopy, radiography can be invasive, expensive, and dangerous, causing life  
139 threatening allergic reactions<sup>5</sup> in some patients, rendering radiography an imperfect  
140 means of SLAP lesion detection<sup>3, 4, 7</sup>.

141           More than 20 provocative tests for the clinical evaluation of SLAP lesions are  
142 proposed in the literature. In most cases, the evaluation of the physical exams by the  
143 original authors reveals promising accuracy for the detection of SLAP lesions<sup>2, 18, 19, 23, 24,</sup>  
144<sup>33, 34</sup>. However, secondary studies often fail to reproduce the initial findings, typically

145 reporting much lower values for the sensitivity and specificity of the physical  
146 examination tests<sup>8, 12, 15, 21, 25, 26</sup>. The discrepancies between the findings most likely  
147 reflect two primary difficulties: 1) the clinical detection of SLAP lesions is hindered by  
148 the fact that SLAP lesions are rarely isolated; meaning they are frequently accompanied  
149 by other various glenohumeral pathologies which are potential sources for labral  
150 symptoms<sup>3</sup>, and 2) differences in study protocols and problems associated with the  
151 methods used to verify accuracy of the physical examinations make comparisons between  
152 studies difficult<sup>8</sup>.

153         The bulk of the literature assesses SLAP lesion tests by determining diagnostic  
154 accuracy through a single verification method. Typically a patient with a suspected  
155 SLAP lesion performs the provocative tests of interest in a clinical setting before  
156 shoulder arthroscopy. The outcome of the SLAP lesion test is then verified with  
157 conclusive arthroscopic findings<sup>7-9, 12, 15, 21, 22, 26</sup>. The results of these comparative studies  
158 have significant quantitative discrepancies, but a fundamental qualitative conclusion  
159 recurs; no single SLAP lesion test has the sensitivity or specificity to independently  
160 determine the presence or absence of a SLAP lesion<sup>7-9, 25</sup>. Although previous studies  
161 assess the diagnostic accuracy of specific SLAP lesion tests, they do little to explain the  
162 reasons behind their apparent failure and rarely suggest or point to any means of  
163 improving the performance of the tests.

164         Clinically based evaluations of SLAP lesion tests account for the majority of  
165 studies to date; however, studies have also assessed test accuracy by attempting to  
166 validate the fundamental design behind various SLAP lesion tests<sup>11, 16, 27, 32</sup>. Provocative  
167 SLAP lesion tests, by definition, function to provoke labral symptoms (primarily pain) as

168 a positive diagnostic sign, by reenacting one of two injury mechanisms. The first  
169 mechanism elicits active tension in the biceps tendon and is typically associated with an  
170 acute traction trauma to the arm or from the accumulation of microtrauma events over  
171 time from repetitive movements such as overhead throwing. The tensile load produced in  
172 the biceps tendon can pull and damage the superior labrum, the functional link between  
173 the insertion of the biceps tendon and the glenoid rim. The second injury mechanism  
174 produces passive compression of the humeral head and is often associated with a fall to  
175 outstretched arms. The compressive load causes superior humeral head translation within  
176 the glenohumeral joint and can result in a collision between the humerus and labrum,  
177 potentially damaging the soft tissue of the labrum<sup>1</sup>. The ability of proposed SLAP lesion  
178 tests to reenact the injury mechanisms that they were designed to replicate has been  
179 examined from several perspectives including anatomic<sup>11, 16, 27</sup>, kinematic<sup>20</sup>, and  
180 electromyographic<sup>11, 32</sup> methods and results illuminate the importance of design  
181 validation during the development of clinical testing procedures.

182         The purpose of this study was to assess the ability of seven provocative tests to  
183 create active tension in the biceps tendon, by characterizing the behaviors of  
184 glenohumeral joint stabilizing muscles, with particular interest in the long head of the  
185 biceps brachii (LHBB) muscle activation and LHBB muscle selectivity. Tests that elicit  
186 larger activation of the LHBB should serve as better diagnostic indicators for SLAP  
187 lesions. Also the other joint stabilizing muscles were examined to determine individual  
188 muscle contributions during the tests, outlining the ability of each test to selectively  
189 activate the LHBB. Selectively activating the LHBB should reduce diagnostic  
190 complications related to the frequent presence of other confounding pathologies with

191 SLAP lesions.

192 ***Materials and Methods.***

193

194 *Subjects and IRB approval*

195 A cohort of 21 healthy volunteers comprised of 11 females ( $24.7 \pm 6.7$  years,  $168.4$   
196  $\pm 5.3$ cm,  $66.9 \pm 9.1$ kg) and 10 males ( $29.4 \pm 10.6$  years,  $178.1 \pm 6.6$  cm,  $80.0 \pm$   
197  $6.4$ kg) with right arm dominance and no history of shoulder pathology were recruited as  
198 subjects in this study. All procedures were approved by the Institutional Review Board at  
199 Boise State University, and all participants read and signed a statement of informed  
200 consent prior to the start of testing.

201

202 *EMG Apparatus and Subject Preparation and Electrode Placement*

203 Electromyography (EMG) was used to record muscle activity for seven muscles  
204 surrounding the dominant glenohumeral joint including the long and short heads of  
205 biceps brachii (LHBB and SHBB), anterior deltoid (DELTA), pectoralis major (PECT),  
206 latissimus dorsi (LAT), infraspinatus (INFRA), and supraspinatus (SUPRA). Each subject  
207 was instrumented with one 44-gage fine-wire indwelling electrode and six surface bipolar  
208 silver-silver chloride EMG electrodes (Noraxon, USA Inc, Scottsdale, AZ). The surface  
209 electrodes were positioned over the muscle belly and parallel with the orientation of the  
210 muscle fibers as suggested by Cram<sup>6</sup>. An additional surface electrode was placed on the  
211 acromion process of the non-dominant shoulder to serve as a reference. Due to the  
212 location of the SUPRA deep to the trapezius, the indwelling electrode was necessary to  
213 acquire SUPRA activity. Using sterile techniques, an emergency medical technician who  
214 was trained specifically for this task by a medical doctor placed the fine-wire indwelling

215 electrode. EMG data were recorded using the Vicon Nexus Software (Vicon, Los  
216 Angelos, CA) at 1250 Hz using a Noraxon Telemetry 900 EMG system (Noraxon USA,  
217 Inc, Scottsdale, AZ).

218

### 219 *EMG Analysis*

220 The EMG signals were analyzed using custom MATLAB software (The  
221 MathWorks, Inc, Natick, MA). A traditional filtering method was used for the EMG  
222 signals for the LHBB, SHBB, DELT, PECT, and SUPRA. Each signal was smoothed by  
223 implementing a root mean square algorithm in combination with a 20ms forward moving  
224 window average. The signals were normalized to a percentage of effort based on their  
225 respective Maximum Voluntary Isometric Contraction (MVIC) peak EMG signal  
226 amplitudes, ideally resulting in a muscle activation range of zero to 100%. Maximum  
227 muscle activation and muscle selectivity were determined for each muscle during each  
228 test. The raw data for the DELT and LAT were processed by the same method, but the  
229 MVIC peak signals were further examined to ensure that the peak amplitude of the signal  
230 did not overlap with a peak from the heartbeat artifacts.

231 Muscle activation and muscle selectivity were calculated to characterize muscle  
232 behavior during the provocative tests, with particular interest in the LHBB behavior.  
233 Muscle activation was used to determine how effective each SLAP lesion test was at  
234 causing individual muscles to activate and was defined as the mean of the peak muscle  
235 activities elicited during the three normalized trials. The ability of each provocative test  
236 to isolate the LHBB is important for diagnosing SLAP lesions due to its common  
237 association with other shoulder pathologies. Therefore in this study, a ratio indicating the

238 ability of each test to selectively activate each muscle was calculated. The muscle  
 239 selectivity for each test was defined by the ratio of the peak activation of the muscle of  
 240 interest over the sum of peak activations for all seven muscles examined, such that a  
 241 selectivity ratio of 1.0 for any muscle, N, would indicate that muscle N was the only  
 242 active muscle contributing to the EMG signal while the other six muscles remained  
 243 inactive. For each test the general selectivity calculation for muscle N was defined as:

$$244 \quad N\_SelectivityRatio = \frac{A_N}{A_{LHBB} + A_{SHBB} + A_{DELT} + A_{PECT} + A_{LAT} + A_{INFRA} + A_{SUPRA}}$$

245  $N$  is the muscle of interest (LHBB, SHBB, DELT, PECT, LAT, INFRA, or SUPRA)

246  $N\_SelectivityRatio$  is the selectivity ratio of muscle  $N$

247  $A_N$  is the peak muscle activation of muscle  $N$

248

#### 249 *Subject Protocol – MVICs and Provocative Tests*

250 Using established EMG protocols, each subject was asked to perform MVICs for  
 251 each muscle of interest on a Biodex System II Dynamometer (Biodex Medical Systems,  
 252 Shirley, NY). Modifications were made to the MVIC recommendations of Cram,  
 253 Hintermeister, and Rowlands<sup>6, 13, 29</sup> to accommodate for the use of the Biodex system  
 254 [Table I]. For each trial, six seconds of data were recorded to ensure that the entire burst  
 255 of muscle activity was captured during each MVIC. The subjects performed three trials  
 256 for each MVIC and were asked to maximally contract for a count of three seconds.  
 257 Subjects rested for thirty seconds between MVIC trials to avoid fatigue effects. For each  
 258 MVIC, the peak amplitude of the EMG signal among the three trials was used to  
 259 normalize the provocative test data.

260 Similarly seven provocative tests were performed based on the descriptions of the  
261 original authors but with modifications to accommodate for use of the Biodex System.  
262 These tests were chosen based on the findings of a preliminary pilot study that evaluated  
263 clinical tests from relevant literature that were designed to reenact either SLAP lesion  
264 injury mechanism<sup>20</sup>. Again the subject performed three trials for a three second count  
265 for each test, six seconds of data were recorded for each trial, and the subjects rested for  
266 thirty seconds between trials.

267

#### 268 *Provocative Test Descriptions and Study Modifications*

269 The modifications for each MVIC and six of the seven SLAP lesion tests utilized the  
270 Biodex System for the purpose of reducing the influence of variances in muscle behavior  
271 and test performance. Tests requiring static resistance against an applied load maintained  
272 the static subject position through the stationary preset up of the Biodex System. Tests  
273 requiring dynamic resistance to an applied load were controlled by the Biodex System  
274 allowing a constant velocity regardless of the force applied by the subject.

275

#### 276 Active Compression Test (ACPD and ACPU)

277 Active Compression has two positions, palm down (ACPD) and palm up (ACPU),  
278 which vary only by rotation of the arm. The patient is standing with the elbow in full  
279 extension, the shoulder is flexed to 90°, and adducted 10–15° medial to the sagittal  
280 plane. For ACPD, the forearm is fully pronated and the glenohumeral joint is  
281 maximally internally rotated such that the thumb points down. The patient is asked to  
282 resist a uniform downward load applied to their arm by the clinician. For ACPU the

283 initial patient positioning is unchanged except the arm is externally rotated such that  
284 the palm faces up. Again, the patient is asked to resist a uniform downward load  
285 applied by the clinician <sup>24</sup>.

286 For the purpose of analysis and due to the nature of the subject population as a  
287 control group having no history of shoulder pathology and therefore asymptomatic,  
288 this study treated ACPU and ACPD as two independent tests. Both tests were  
289 modified such that the subject was seated in the Biodex. The orientation of the  
290 subject's arm remained true to O'Brien's original description of the test, but the  
291 subject was asked to resist the stationary position of the Biodex arm by attempting to  
292 lift his/her arm superiorly for both ACPD and ACPU.

293

#### 294 Speed's Test (Speed's)

295 According to the original description, the patient is standing and resists a  
296 downward force applied to the upper extremity with the elbow extended, forearm  
297 supinated, and arm elevated to 90° <sup>2</sup>. In this study the orientation of the arm remained  
298 similar to the original definition, but Speed's test was modified into a dynamic  
299 movement controlled by the Biodex System. The subject's arm started hanging  
300 beside and parallel to the body, and then the subject was asked to raise the arm (flex  
301 the shoulder) with as much force as possible to 90° . Regardless of the force applied  
302 by the subject, motion was restricted to a constant velocity by the Biodex System of  
303 60° per second.

304

#### 305 Pronated Load Test (ProLoad)

306 In the original description, the patient is in the seated position with elbow flexed  
307 to 90°, the arm is abducted to 90°, maximally externally rotated, and the forearm is  
308 fully pronated. The patient is then asked to perform an isometric contraction of the  
309 Biceps<sup>33</sup>. The Pronated Load Test was negligibly modified for this study. The  
310 subject sat in the original orientation in the Biodex System, which was set up such  
311 that the arm was supported just proximal to the elbow. The subject was asked to  
312 perform a bicep contraction (pronated curl) which was resisted by the static set up of  
313 the Biodex System.

314

#### 315 Biceps Load I Test (Bicep I)

316 The patient is in the supine position when an anterior apprehension test is  
317 performed starting with the arm abducted 90° and the forearm fully supinated  
318 according to its original definition<sup>19</sup>. Bicep I was modified such that the patient was  
319 seated in the same position as ProLoad, except the forearm was fully supinated. The  
320 subject was asked to perform a bicep contraction (curl), which was resisted by the  
321 static set up by the Biodex System.

322

#### 323 Biceps Load II Test (Bicep II)

324 The patient is supine with the arm abducted to 120° degrees, the elbow flexed to  
325 90° degrees, and the forearm fully supinated. The patient is then asked to flex the  
326 elbow against the resistance of the clinician<sup>18</sup>. For this study the modification for  
327 Bicep II paralleled those made to Bicep I except the arm was abducted to  
328 120° degrees instead of 90°.

329

330 Resisted Supination External Rotation Test (RSER)

331 The authors describe putting the patient in the supine position with scapula near  
332 the edge of the table, the patients arm is supported by the physician at the wrist, with  
333 the arm is abducted to 90° and the elbow is flexed between 65° and 70° degrees. The  
334 clinician then externally rotates the arm while the patient is asked to supinate the  
335 forearm<sup>23</sup>. The RSER test was essentially unchanged for this study, and the  
336 movement was not controlled by the Biodex System. One board certified athletic  
337 trainer performed RSER with each subject in the study in the supine position on the  
338 Biodex.

339

340 Supination Sign Test (Yergason's)

341 In the seated position with the elbow flexed to 90° and forearm fully pronated,  
342 the patient is asked to attempt supination of the forearm while the physician resists  
343 the motion while holding the wrist<sup>34</sup>. Yergason's was scarcely modified; as the  
344 patient maintained the defined orientation but with the forearm fastened to the static  
345 Biodex arm. The patient was asked to attempt to supinate the forearm against the  
346 static setup of the Biodex.

347

348 *Statistical Analysis*

349 Statistical analysis was conducted using SPSS Statistics Software (SPSS, Inc,  
350 Chicago, IL) to determine significant differences in maximum muscle activations and  
351 muscle selectivities for each test, between tests, and between male and female groups. A

352 repeated measures analysis of variance (ANOVA) test was performed to identify  
353 significant differences between provocative tests for each individual muscle. A pair-wise  
354 T-test post-hoc analysis was performed to compare results between each test using a p-  
355 value sliding scale Bonferroni adjustment <sup>14</sup>. Likewise, a paired-sample T-test was used  
356 to examine potential differences in muscle activation ( $p = 0.05$ ) between male and female  
357 groups.

358 **Results.**

359 A post-hoc pair-wise comparison between males and females showed no  
360 differences between male and female groups for any muscle or for any provocative test  
361 with all p-values exceeding 0.05. Therefore male and female data were pooled for all  
362 subsequent statistical analyses. For each individual muscle, the repeated measures  
363 ANOVA analysis found significant differences in both muscle activation and muscle  
364 selectivity among the eight provocative tests ( $p < .05$ ).

365 To determine which provocative tests resulted in the greatest activations for the  
366 individual muscles, 28 pair-wise comparisons between the eight tests were made for each  
367 muscle. Each muscle analyzed showed a significant difference in peak muscle activity  
368 between one or more of the pairs of provocative tests with the exception of the LAT.  
369 Specifically, the LHBB demonstrated a significant difference ( $p=.000$ ) in activity  
370 between tests. The eight statistically significant pair-wise comparisons enabled the tests  
371 to be characterized into one of two performance groups based on their respective LHBB  
372 activation; high performing and low performing. Speed's, ACPU, Bicep I, and Bicep II,  
373 tests were 'high performing', eliciting the largest mean peak EMG amplitudes without  
374 statistical differences among the four tests, while RSER, Yergason's, ACPD, and  
375 ProLoad were classified as 'low performing' (Figure 2). The mean normalized peak  
376 activations (% MVIC) for each muscle elicited during all eight tests are noted in Table II.

377 The statistical analysis with regards to muscle selectivity for each test proved  
378 similar to those for muscle activation. There were significant differences in muscle  
379 selectivity across the provocative tests ( $p=.000$ ). A post-hoc pair-wise comparison  
380 showed that one or more pairs of tests had significant differences in muscle selectivity for

381 each muscle with the exception of the LAT and INFRA. The eleven statistically  
382 significant pairs allowed the tests to be categorized into high and low performance groups  
383 based on LHBB selectivity. RSER, Bicep I, Bicep II, and Yergason's tests were 'high  
384 performing', recruiting the LHBB more selectively than ProLoad, Speed's, ACPU, and  
385 ACPD, which were categorized as 'low performing' (Figure 3). Again there was no  
386 statistical difference among tests within each group. The mean selectivities of each  
387 muscle for all eight tests are noted in Table III.  
388

389 *Discussion.*

390           The aim of this study was to characterize the muscle behavior of seven  
391 glenohumeral joint stabilizing muscles, focusing on the LHBB, during eight modified  
392 provocative tests that were designed to detect SLAP lesions by loading the biceps tendon  
393 in tension through LHBB activation. The active tension in the biceps tendon is thought to  
394 reproduce the injury mechanism of a SLAP lesion, which should provoke a response  
395 from suspected SLAP lesion patients yielding a positive diagnostic sign<sup>1,33</sup>. In this study,  
396 Bicep I and Bicep II were the most promising SLAP lesion tests according to their  
397 favorable LHBB behavior, eliciting high LHBB activity while remaining highly selective  
398 for the LHBB, indicating these two tests should function effectively as assessment tools  
399 for the clinical evaluation of SLAP lesions.

400           The magnitude of LHBB activation during each of the clinical evaluations is a  
401 measure of the sensitivity of the maneuver to incite active tension in the LHBB tendon  
402 which should increase the likelihood of detecting a SLAP tear. Although EMG signal  
403 amplitude cannot be directly related to muscle force in most cases, the tests that most  
404 strongly activate LHBB should provide relatively higher traction forces to the superior  
405 labrum. Speed's, ACPU, Bicep I, and Bicep II tests produced the largest LHBB activities,  
406 reaching above 90 % MVIC, suggesting that a greater respective load was applied to the  
407 biceps tendon during these tests. Although none of the tests apply loads sufficient to  
408 produce a SLAP lesion, Speed's ACPU, Bicep I, and Bicep II tests created the largest  
409 LHBB activation and therefore reproduced the injury mechanism more effectively than  
410 the other four low-performing tests (RSER, Yergason's, ACPD, and ProLoad). Although

411 SLAP lesion test assessment is prevalent in the literature, comparison between studies is  
412 difficult due to the lack of overlap of tests between similar studies. However, two studies  
413 support the findings of this study in that ACPU and Bicep II have both been reported to  
414 elicit large LHBB EMG amplitudes<sup>11 32</sup>.

415 LHBB selectivity served as an equally important variable to consider for  
416 characterizing LHBB behavior and for assessing SLAP lesion tests, as it is an indicator of  
417 test specificity. The diagnostic accuracy of SLAP lesion tests are often hindered by the  
418 frequent occurrence of other glenohumeral pathologies, such as rotator cuff tears, that  
419 make determining the origin of shoulder symptom challenging at best<sup>9, 17, 21, 31</sup>.  
420 Consequently provocative tests that are able to isolate the LHBB would be beneficial  
421 because high LHBB selectivity denotes a lesser contribution from other joint stabilizing  
422 muscles that can produce a false SLAP lesion diagnosis. RSER, Bicep I, Bicep II, and  
423 Yergason's tests were 'high performing' with regards to selectively recruiting the LHBB.  
424 Each high performing test resulted in LHBB selectivity between 0.23 and 0.25, compared  
425 to the range of 0.12 and 0.16 selectivity for the 'low performing' tests (Proload, Speeds,  
426 ACPU, and ACPD). Unfortunately LHBB selectivity is not reported elsewhere in the  
427 relevant literature, but these results concur with the findings of the preliminary pilot study  
428 <sup>20</sup>.

429 The two overall top performing SLAP lesion tests, Bicep I and Bicep II, elicited  
430 large LHBB activation while demonstrating high LHBB selectivity. The clinical  
431 implications derived from the remaining tests that were 'high performing' in only a single  
432 area of LHBB behavior, either highly specific (activation – ACPU and Speed's) or highly  
433 sensitive (selective – RSER and Yergason's), may be limited if used on their own. Top

434 performing SLAP lesion tests, that elicited large LHBB activation and were highly  
435 selective for the LHBB, should be closely examined in hopes of defining the  
436 characteristics that may be responsible for their promising LHBB behavior.

437         Bicep I and Bicep II are very similar tests, varying only by the flexion of the  
438 shoulder joint. Bicep I, Bicep II, Speed's, and ACPU, all of have desirable behavior in  
439 one or both suites and may be useful for future work, by examining the clinical  
440 implications of these tests in combination. These four tests, Bicep I, Bicep II, Speed's,  
441 and ACPU, share similar test and design characteristics relating to location of the applied  
442 load, forearm orientation, joint position, and line of pull during either a static or dynamic  
443 provocative test designed to activate the LHBB. Each of these tests was performed with  
444 a supinated forearm and required active resistance to an external load applied  
445 perpendicular to the palm of the subject's hand. Each high performing test was  
446 performed in one of two joint positions which placed the LHBB and biceps tendon in a  
447 direct line of pull with the superior labrum. The first joint position (Speed's and ACPU)  
448 flexed the shoulder to a maximum of 90° with the elbow fully extended. The second  
449 joint position (Bicep I and Bicep II) had the shoulder abducted at or above 90° with the  
450 elbow flexed at 90°. The major difference between these four tests is the way the tests  
451 are performed; Speed's is a dynamic test while Bicep I, Bicep II, and ACPU are static  
452 tests, where the patient resists the load without the ability to move.

453         In this study ACPU and the Speed's were extremely similar and although both  
454 were 'high performing' for LHBB activation, their differences may prove important  
455 means of understanding the role and importance of SLAP lesions test characteristics. The  
456 tests have slight differences in patient orientation and type of movement; ACPU places

457 the arm medial to the sagittal plane and the test is static, while Speed's is parallel to the  
458 medial plane and involves a dynamic movement. These small differences may have  
459 important consequences, and a close examination of these kinds of test characteristics and  
460 their relation to test performance may help illuminate a means of improving test design  
461 and accuracy.

462         Although the focus of this study was the behavior of the LHBB, six other joint  
463 stabilizing muscles were recorded to enable LHBB selectivity calculations and in hopes  
464 of characterizing any other muscle behaviors or patterns. Peak muscle activities and  
465 muscle selectivity were examined for all remaining muscles (SHBB, DELT, PECT, LAT,  
466 INFRA, and SUPRA), and statistical analysis revealed that it may be unnecessary to  
467 monitor the LAT and INFRA during these tests, because none of the tests had a  
468 significant difference in terms of activation of the LAT or in selectively isolating either  
469 the LAT or INFRA muscles.

470         The primary inherent limitation of this study is that the subjects had no history of  
471 shoulder pathology; therefore labral symptoms were not used as a means to assess SLAP  
472 test performance. Also the healthy subject pool may misrepresent SLAP lesion patients  
473 due to the potential for differences in muscle behavior between healthy subjects and those  
474 with labral pathology. Furthermore, the EMG signals were all normalized based on peak  
475 activities elicited during MVIC, and results exceeded 100% in some cases and may make  
476 comparison between subjects difficult. Specifically, the dynamic Speed's test, which had  
477 the largest mean activation (140.9% MVIC) among the tests, was not a surprising finding,  
478 as the dynamic movement was normalized to a static MVIC. Muscle activation is known  
479 to vary with both muscle length and shortening or lengthening velocity. Therefore,

480 comparing activation during a dynamic test to data collected in a static configuration may  
481 not be optimal. For the static tests, LHBB activations were generally below or much less  
482 than that of Speed's, suggesting that the normalization procedure was more appropriate  
483 for those tests. However, in some tests subjects were able to achieve more than 100%  
484 MVIC in some muscles, which means either that the tests were more effective in isolating  
485 those muscles than the MVIC configurations, or that slight differences in positioning or  
486 in subject effort in the clinical tests and the MVIC tests affected the muscle activation  
487 values recorded.

488 Future studies would improve on the scope of this study by recruiting subjects  
489 who have a suspected SLAP lesion and are scheduled for arthroscopic assessment.  
490 Employing the methods and results of this study, improvements would utilize the  
491 promising LHBB behavior of the top performing modified tests (Bicep I and Bicep II) in  
492 conjunction with analyses of associated joint torques. Although joint torque data was  
493 not collected in this study due to the inability to acquire torque information for all of the  
494 eight modified tests, the top performing SLAP lesions tests are oriented such that the  
495 Biodex System could easily provide such information. An analysis of joint torques and  
496 associated loads during these tests may further quantify the ability of these tests to create  
497 tension in the biceps tendon.

498 Recent studies utilizing arthroscopic verification for clinical evaluations have  
499 documented a drastic increase in SLAP lesion detection by using the indications of two or  
500 more SLAP tests, specifically when at least one test is highly sensitive and another is  
501 highly specific<sup>8,25</sup>. Consequently, assessing the array of 'high performing' test

502 combinations, utilizing various combinations of single suite high performance tests with  
503 various test characteristics may have surprising results and prove worthwhile.

504           Lastly, although difficult to determine and requiring a large pool of control and  
505 experimental data, comparisons between the muscle behaviors of a healthy population  
506 and those who have a suspected SLAP lesion may illuminate some general pattern  
507 differences that could be indicative of SLAP lesions and be useful for furthering clinical  
508 diagnostic techniques and accuracy.

509

510 *Conclusions.*

511 In summary, modified versions of Bicep I and Bicep II resulted in the greatest  
512 LHBB activation and LHBB selectivity of the SLAP lesion tests in this study. ACPU,  
513 and Speed's resulted in the large LHBB activation, but were not selective for the LHBB.  
514 Bicep I, Bicep II, ACPU, and Speed's each elicit some promising LHBB behavior, and  
515 maybe useful in combination to aid the clinical detection of SLAP lesions. These four  
516 tests utilize a unique range of test variables that may prove valuable for optimal SLAP  
517 test design and function. Future studies should evaluate the importance of these  
518 variables, incorporate joint torque analyses, and expand the scope of the study to include  
519 patients who have a suspected SLAP lesion to optimize, validate, and improve the  
520 diagnostic accuracy of provocative SLAP lesion test.

521 *Acknowledgements*

522 This project was supported by grant number 1-R15 AR054637 through the  
523 National Institute of Arthritis and Musculoskeletal and Skin Diseases. Special thanks to  
524 Intermountain Orthopedics with particular thanks to Drs. Kurt J. Nilsson, MD, and Kevin  
525 G. Shea, MD for their support of this study.

526

527

**References.**

528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573

1. Andrews, J., W. Carson, Jr, and W. McLeod, *Glenoid labrum tears related to the long head of the biceps*. Am J Sports Med, 1985. **13**(5): p. 337-341.
2. Bennett, W.F., *Specificity of the Speed's Test: Arthroscopic Technique for Evaluating the Biceps Tendon at the Level of the Bicipital Groove*. The Journal of Arthroscopic and Related Surgery, 1998. **14**(8): p. 789-796.
3. Chandnani, V., et al., *Glenoid labral tears: prospective evaluation with MRI imaging, MR arthrography, and CT arthrography*. Am. J. Roentgenol., 1993. **161**(6): p. 1229-1235.
4. Chang, D., et al., *SLAP lesions: Anatomy, clinical presentation, MR imaging diagnosis and characterization*. European Journal of Radiology, 2008. **68**(1): p. 72-87.
5. Chundru, U., *Magnetic Resonance Arthrography*. Musculoskeletal, 2009.
6. Cram, J.R. and G.S. Kasman, *Introduction to surface electromyography*. 1998, Gaithersburg, MD: Aspen Publishers Inc.
7. DaSilva, J.J., A.G. McNickle, and B.J. Cole, *Arthroscopic Superior Labrum Anterior to Posterior Repair*. Operative Techniques in Orthopaedics, 2008. **18**(1): p. 53-61.
8. Dessaur, W. and M. Magarey, *Diagnostic accuracy of clinical tests for superior labral anterior posterior lesions: a systematic review*. Journal of Orthopaedic and Sports Physical Therapy, 2008. **38**(6): p. 341-352.
9. Gill, H.S., et al., *Physical Examination for Partial Tears of the Biceps Tendon*. Am J Sports Med, 2007. **35**(8): p. 1334-1340.
10. Golay, S.A.M., 'Smoothing and differentiation of data, by simplified least squares procedures'. Analytical Chemistry, 1964. **36**: p. 1627-1639.
11. Green, R.A., et al., *An evaluation of the anatomic basis of the O'Brien active compression test for superior labral anterior and posterior (SLAP) lesions*. Journal of Shoulder and Elbow Surgery, 2008. **17**(1): p. 165-171.
12. Guanche, C.A. and D.C. Jones, *Clinical Testing for Tears of the Glenoid Labrum*. The Journal of Arthroscopic and Related Surgery, 2003. **19**(5): p. 517-523.
13. Hintermeister, R.A., et al., *Electromyographic Activity and Applied Load During Shoulder Rehabilitation Exercises Using Elastic Resistance*. The American Journal of Sports Medicine, 1998. **26**(2): p. 210-220.
14. Holm, S., *A simple sequentially rejective multiple test procedure*. Scandinavian Journal of Statistics, 1979. **6**: p. 65-70.
15. Holtby, R. and H. Razmjou, *Accuracy of the Speed's and Yergason's Tests in Detecting Biceps Pathology and SLAP Lesions: Comparison With Arthroscopic Findings*. The Journal of Arthroscopic and Related Surgery, 2004. **20**(3): p. 231-236.

- 574 16. Keren J. Shanley, R.A.G., Nicholas F. Taylor., *An evaluation of the anatomical*  
575 *basis of the Biceps Load tests I and II for superior labral anterior and posterior*  
576 *(SLAP) lesions*. Clinical Anatomy, 2008. **21**(7): p. 647-651.
- 577 17. Kibler, W.B., *Specificity and Sensitivity of the Anterior Slide Test in Throwing*  
578 *Athletes With Superior Glenoid Labral Tears*. Arthroscopy, 1995. **11**(3): p. 296-  
579 300.
- 580 18. Kim, S.-H., et al., *Biceps Load Test II: A Clinical Test for SLAP Lesions of the*  
581 *Shoulder*. The Journal of Arthroscopic and Related Surgery, 2001. **17**(2): p. 160-  
582 164.
- 583 19. Kim, S.-H., K.-I. Ha, and K.-Y. Han, *Biceps Load Test: A Clinical Test for*  
584 *Superior Labrum Anterior and Posterior Lesions in Shoulders With Recurrent*  
585 *Anterior Dislocations*. The American Journal of Sports Medicine, 1999. **27**(3): p.  
586 300-303.
- 587 20. Kuhlman, S., Sabick, M, Pfeirffer., *Biomechanical Evaluation of Physical*  
588 *Examination Tests for Superior Labrum Anterior to Posterior (SLAP) Lesions*.  
589 Pending Review, 2009.
- 590 21. McFarland, E.G., T.K. Kim, and R.M. Savino, *Clinical Assessment of Three*  
591 *Common Tests for Superior Labral Anterior-Posterior Lesions*. Am J Sports Med,  
592 2002. **30**(6): p. 810-815.
- 593 22. Munro, W. and R. Healy, *The validity and accuracy of clinical tests used to detect*  
594 *labral pathology of the shoulder* “ A systematic review. Manual Therapy. **In**  
595 **Press, Corrected Proof**.
- 596 23. Myers, T.H., J.R. Zemanovic, and J.R. Andrews, *The Resisted Supination*  
597 *External Rotation Test: A New Test for the Diagnosis of Superior Labral Anterior*  
598 *Posterior Lesions*. The American Journal of Sports Medicine, 2005. **33**(9): p.  
599 1315-1320.
- 600 24. O'Brien, S.J., et al., *The Active Compression Test: A New and Effective Test for*  
601 *Diagnosing Labral Tears and Acromioclavicular Joint Abnormality*. The  
602 American Journal of Sports Medicine, 1998. **26**(5): p. 610-613.
- 603 25. Oh, J.H., et al., *The Evaluation of Various Physical Examinations for the*  
604 *Diagnosis of Type II Superior Labrum Anterior and Posterior Lesion*. Am J  
605 Sports Med, 2008. **36**(2): p. 353-359.
- 606 26. Parentis, M.A., et al., *An Evaluation of the Provocative Tests for Superior Labral*  
607 *Anterior Posterior Tears*. Am J Sports Med, 2006. **34**(2): p. 265-268.
- 608 27. Parentis, M.A., et al., *An Anatomic Evaluation of the Active Compression Test*.  
609 Journal of Shoulder and Elbow Surgery, 2004. **13**(4): p. 410-416.
- 610 28. Robert, D., *Savitsky-Golay Filters*. 2008.
- 611 29. Rowlands, L., et al., *Kinesiology of the Empty Can Test*. Am J Phys Med Rehabil,  
612 1995. **74**(4): p. 302-304.
- 613 30. Snyder, S.J., et al., *SLAP Lesions of the Shoulder*. The Journal of Arthroscopic  
614 and Related Surgery, 1990. **6**(4): p. 274-279.
- 615 31. Stetson, W.B. and K. Templin, *The Crank Test, the O'Brien Test, and Routine*  
616 *Magnetic Resonance Imaging Scans in the Diagnosis of Labral Tears*. Am J  
617 Sports Med, 2002. **30**(6): p. 806-809.

- 618 32. Swearingen, J.C., et al., *Electromyographic analysis of physical examination tests*  
619 *for type II superior labrum anterior-posterior lesions*. Journal of Shoulder and  
620 Elbow Surgery, 2006. **15**(5): p. 576-579.
- 621 33. Wilk, K.E., et al., *Current Concepts in the Recognition and Treatment of Superior*  
622 *Labral (SLAP) Lesions*. Journal of Orthopaedic and Sports Physical Therapy,  
623 2005. **35**: p. 273-291.
- 624 34. Yergason, R.M., *Supination Sign*. 1995.  
625  
626

627 **Tables**

628 **Table I:** MVIC joint positions and resisted maneuvers for the seven muscles of interest.

629 **Table II:** Resulting mean normalized peak muscle activations (%MVIC) and standard

630 deviations monitored during each SLAP lesion test.

631 **Table III:** Resulting mean muscle selectivity values and standard deviations monitored

632 during each SLAP lesion test.

633

634

<b>Muscle</b>	<b>Joint Position</b>	<b>Resisted Maneuver</b>
<b>DELTA</b>	Arm at side	Shoulder flexion
<b>LHBB / SHBB</b>	Elbow flexed 90°, shoulder flexed 90°	Elbow flexion
<b>INFRA</b>	Arm abducted 45°, elbow flexed 90°	External Rotation
<b>LAT</b>	Shoulder flexed 90°, arm internally rotated	Shoulder extension
<b>PECT</b>	Arm abducted 90°, forearm supinated	Horizontal adduction
<b>SUPRA</b>	Arm abducted 90°, forward flexed 30°, and internally rotated	Maintain against resistance

635

Muscle Peak Mean Activation (% MVIC) and Standard Deviation During SLAP Lesion Tests														
	LHBB		SHBB		DELTA		PECT		LAT		INFRA		SUPRA	
ACPD	116.6	(75.2)	86.7	(60.4)	192.1	(97.6)	89.2	(42.8)	116.2	(161.6)	182.3	(174.8)	84.8	(38.6)
ACPU	74.9	(66.4)	15.7	(12.8)	116.7	(65.7)	28.4	(17.5)	93.8	(146.3)	110.1	(82.9)	105.4	(62.7)
Speeds	140.9	(100.9)	104.2	(90.6)	158.4	(72.0)	88.0	(35.4)	124.1	(158.4)	143.3	(96.4)	107.7	(58.5)
Bicep I	97.6	(37.2)	88.9	(36.5)	43.9	(49.4)	36.0	(20.0)	72.6	(65.4)	56.8	(58.9)	26.7	(22.5)
Bicep II	94.0	(48.0)	88.7	(42.8)	49.8	(56.9)	44.8	(21.8)	56.2	(46.4)	41.9	(37.9)	30.7	(43.1)
ProLoad	58.1	(32.8)	39.8	(19.0)	58.9	(49.5)	28.5	(15.9)	70.1	(60.4)	69.2	(53.7)	39.4	(31.1)
RSER	89.2	(65.7)	85.6	(65.0)	15.4	(17.4)	23.5	(19.5)	49.5	(32.6)	59.2	(61.1)	25.9	(22.5)
Yergasons	81.1	(46.3)	81.6	(52.0)	22.3	(19.1)	31.8	(19.9)	98.4	(152.8)	56.3	(50.4)	24.1	(19.7)

636

637

Muscle Mean Selectivity and Standard Deviation During SLAP Lesion Tests														
	LHBB		SHBB		DELTA		PECT		LAT		INFRA		SUPRA	
ACPD	0.132	(0.056)	0.118	(0.059)	0.300	(0.104)	0.225	(0.139)	0.268	(0.203)	0.633	(0.145)	0.105	(0.049)
ACPU	0.122	(0.091)	0.034	(0.032)	0.259	(0.118)	0.102	(0.074)	0.264	(0.178)	0.494	(0.200)	0.213	(0.122)
Speeds	0.152	(0.065)	0.138	(0.066)	0.263	(0.076)	0.209	(0.076)	0.283	(0.214)	0.566	(0.182)	0.131	(0.064)
Bicep I	0.244	(0.079)	0.308	(0.124)	0.188	(0.134)	0.229	(0.110)	0.479	(0.206)	0.587	(0.230)	0.064	(0.055)
Bicep II	0.231	(0.070)	0.303	(0.113)	0.217	(0.135)	0.293	(0.146)	0.452	(0.188)	0.558	(0.281)	0.072	(0.082)
ProLoad	0.160	(0.069)	0.144	(0.078)	0.204	(0.124)	0.150	(0.078)	0.401	(0.252)	0.602	(0.220)	0.113	(0.087)
RSER	0.255	(0.086)	0.336	(0.136)	0.092	(0.080)	0.164	(0.106)	0.415	(0.229)	0.636	(0.203)	0.075	(0.047)
Yergasons	0.225	(0.086)	0.311	(0.158)	0.104	(0.063)	0.186	(0.086)	0.427	(0.226)	0.653	(0.189)	0.067	(0.044)

638

639


640 **Illustrations and Legends**

641 **Figure 1:** Example of Biodex System modifications, ACPD and ACPU.

642 **Figure 2:** LHBB mean muscle activation (%MVIC) for each SLAP test.


643 **Figure 3:** LHBB mean muscle selectivity for each SLAP test.

644




645  
646  
647  
648  
650  
651  
652  
653  
654  
655  
656

657



658



659