

2-2023

The Relationship Between Fundamental Motor Skills and Physical Fitness in Children Aged 7-12 Years

Caifang Wu
Southern Connecticut State University

Xiaozan Wang
East China Normal University

Qinping Xu
East China Normal University

Lingsong Kong
University of Massachusetts Amherst

Follow this and additional works at: <https://scholarworks.boisestate.edu/ijpah>



Part of the [Exercise Science Commons](#), [Health and Physical Education Commons](#), [Public Health Commons](#), and the [Sports Studies Commons](#)

Recommended Citation

Wu, Caifang; Wang, Xiaozan; Xu, Qinping; and Kong, Lingsong (2023) "The Relationship Between Fundamental Motor Skills and Physical Fitness in Children Aged 7-12 Years," *International Journal of Physical Activity and Health*: Vol. 2: Iss. 1, Article 26.

DOI: <https://doi.org/10.18122/ijpah.020126.boisestate>

Available at: <https://scholarworks.boisestate.edu/ijpah/vol2/iss1/26>

The Relationship Between Fundamental Motor Skills and Physical Fitness in Children Aged 7-12 Years

Abstract

The purpose of this study was to examine the relationship between fundamental motor skills (FMS) and physical fitness (PF) in children aged 7-12 years. The participants were 217 children (50.7% female) from a primary school in Shanghai. The FMS was assessed using Chinese version Test of Gross Motor Development–3 edition (CTGMD-3), which includes 6 locomotor skills and 7 ball skills. PF was measured following China's national physical health standards for students, which includes BMI, vital capacity, sit and reach, 50m sprint, one-minute rope jumping, one-minute sit-ups and 50m * 8 shuttle run. The weighted score of the 7 tests was calculated to represent the performance of PF. Four participants were assigned into one group and took all tests within a physical education class. Two well-trained graduate assistants performed the tests within one month. Descriptive statistics were computed after checking the normality of the scores of CTGMD-3 and PF. Pearson correlation was used to determine the relationship between FMS and PF. The significance level was set at 0.05. The results showed a significantly positive correlation between the CTGMD-3 score and PF score ($r = 0.166, p = 0.014$). There was a significantly positive correlation between the score of locomotor skills and PF scores ($r = 0.269, p < 0.01$). No significant correlation between the score of ball skills and PF score ($r = 0.065, p = 0.338$) was found. It is concluded that FMS, especially locomotor skills, was associated with PF among children aged 7-12 years. The generalizability of this study was limited since the participants were from one primary school in Shanghai. The small sample size might limit the power of this study. Future research is needed to explore the mechanism underlying the relationship between FMS and PF.

The Relationship Between Fundamental Motor Skills and Physical Fitness in Children Aged 7-12 Years

Caifang Wu^a, Xiaozan Wang^b, Qiping Xu^b, and Lingsong Kong^c

^aSouthern Connecticut State University, ^bEast China Normal University, College of Physical Education and Health, ^cUniversity of Massachusetts Amherst

Abstract

The purpose of this study was to examine the relationship between fundamental motor skills (FMS) and physical fitness (PF) in children aged 7-12 years. The participants were 217 children (50.7% female) from a primary school in Shanghai. The FMS was assessed using Chinese version Test of Gross Motor Development–3 edition (CTGMD-3), which includes 6 locomotor skills and 7 ball skills. PF was measured following China's national physical health standards for students, which includes BMI, vital capacity, sit and reach, 50m sprint, one-minute rope jumping, one-minute sit-ups and 50m * 8 shuttle run. The weighted score of the 7 tests was calculated to represent the performance of PF. Four participants were assigned into one group and took all tests within a physical education class. Two well-trained graduate assistants performed the tests within one month. Descriptive statistics were computed after checking the normality of the scores of CTGMD-3 and PF. Pearson correlation was used to determine the relationship between FMS and PF. The significance level was set at 0.05. The results showed a significantly positive correlation between the CTGMD-3 score and PF score ($r = 0.166, p = 0.014$). There was a significantly positive correlation between the score of locomotor skills and PF scores ($r = 0.269, p < 0.01$). No significant correlation between the score of ball skills and PF score ($r = 0.065, p = 0.338$) was found. It is concluded that FMS, especially locomotor skills, was associated with PF among children aged 7-12 years. The generalizability of this study was limited since the participants were from one primary school in Shanghai. The small sample size might limit the power of this study. Future research is needed to explore the mechanism underlying the relationship between FMS and PF.

Keywords: fundamental motor skills, physical fitness, BMI, locomotor skills