

**CLASSIFICATION OF VERTEX-TRANSITIVE
STRUCTURES**

by
Stephanie Potter

A thesis
submitted in partial fulfillment
of the requirements for the degree of
Master of Science in Mathematics
Boise State University

May 2017

© 2017
Stephanie Potter
ALL RIGHTS RESERVED

BOISE STATE UNIVERSITY GRADUATE COLLEGE

DEFENSE COMMITTEE AND FINAL READING APPROVALS

of the thesis submitted by

Stephanie Potter

Dissertation Title: Classification of Vertex-Transitive Structures

Date of Final Oral Examination: 14 March 2017

The following individuals read and discussed the dissertation submitted by student Stephanie Potter, and they evaluated her presentation and response to questions during the final oral examination. They found that the student passed the final oral examination.

Samuel Coskey, Ph.D. Chair, Supervisory Committee

John Clemens, Ph.D. Member, Supervisory Committee

Marion Scheepers, Ph.D. Member, Supervisory Committee

The final reading approval of the thesis was granted by Samuel Coskey, Ph.D., Chair of the Supervisory Committee. The thesis was approved by the Graduate College.

ABSTRACT

When one thinks of objects with a significant level of symmetry it is natural to expect there to be a simple classification. However, this leads to an interesting problem in that research has revealed the existence of highly symmetric objects which are very complex when considered within the framework of Borel complexity. The tension between these two seemingly contradictory notions leads to a wealth of natural questions which have yet to be answered.

Borel complexity theory is an area of logic where the relative complexities of classification problems are studied. Within this theory, we regard a classification problem as an equivalence relation on a Polish space. An example of such is the isomorphism relation on the class of countable groups. The notion of a Borel reduction allows one to compare complexities of various classification problems.

The central aim of this research is to determine the Borel complexities of various classes of vertex-transitive structures, or structures for which every pair of elements are equivalent under some element of its automorphism group. John Clemens has shown that the class of vertex-transitive graphs has maximum possible complexity, namely Borel completeness. On the other hand, we show that the class of vertex-transitive linear orderings does not.

We explore this phenomenon further by considering other natural classes of vertex-transitive structures such as tournaments and partial orderings. In doing so, we discover that several other complexities arise for classes of vertex-transitive structures.

TABLE OF CONTENTS

ABSTRACT	iv
LIST OF FIGURES	vi
1 Introduction	1
2 Vertex-Transitive Graphs and Partial Orders	8
3 Vertex-Transitive Linear Orderings	19
4 Vertex-Transitive Tournaments	29
REFERENCES	36

LIST OF FIGURES

1.1	Hierarchy of relevant benchmark equivalence relations	3
1.2	Structures studied in this thesis	5
2.1	Depiction of $u_i R u_j$ with $u_i = g_p N$ and $u_j = g_q N$	11
2.2	Graph of M as constructed by Mekler, [8]	13
2.3	Resulting square if there exist $k_1, k_2 \in \mathbb{N}$ with $v_{k_1} \text{ adj } v_{k_2} \in G$	15
4.1	Directed edges from vertex $(0, 0)$ to vertices one column to the right. . .	32

CHAPTER 1

INTRODUCTION

Classification problems in Mathematics ask the question: “How can objects of a given type be identified and distinguished from one another, up to some equivalence relation?”

When one thinks of objects with a significant level of symmetry it is natural to expect there to be a simple classification. However, this leads to an interesting problem in that research has revealed the existence of highly symmetric objects which are very complex when considered within the framework of Borel complexity. The tension between these two seemingly contradictory notions leads to a wealth of natural questions which have yet to be answered. In this thesis we will answer some of these questions in an attempt to further understand where vertex-transitive structures lie in terms of Borel complexity theory.

Borel complexity theory is an area of logic where the relative complexities of classification problems are studied. Within this theory, we regard a classification problem as an equivalence relation on a Polish space. An example of such is the isomorphism relation on the class of countable groups.

More generally, we can instead consider any invariant Borel class of countable structures, which we define below.

Definition 1. Let $L = \{R_i : i \leq I\}$ be a finite relational language, where R_i has arity n_i . Denote the **space of all countable L -models** as $\text{Mod}(L)$. Each element of $\text{Mod}(L)$ can be viewed as an element of the product space

$$X_L = \prod_{i \leq I} 2^{\mathbb{N}^{n_i}}.$$

That is, for every $x \in X_L$ let $M_x \in \text{Mod}(L)$ be the countable model coded by x . Then for any $i \in I$ and $(k_1, \dots, k_{n_i}) \in \mathbb{N}^{n_i}$, it is the case that $R_i^{M_x}(k_1, \dots, k_{n_i}) \Leftrightarrow x_i(k_1, \dots, k_{n_i}) = 1$. For the remainder of this paper we will identify $\text{Mod}(L)$ and X_L .

Definition 2. The **logic action** of S_∞ on $\text{Mod}(L)$ is defined by letting $g \cdot M = N$ if and only if

$$R_i^N(k_1, \dots, k_{n_i}) \Leftrightarrow R_i^M(g^{-1}(k_1), \dots, g^{-1}(k_{n_i}))$$

for all $i \in I$ and $(k_1, \dots, k_{n_i}) \in \mathbb{N}^{n_i}$. Therefore, $g \cdot M = N$ if and only if g is an isomorphism from M onto N .

Definition 3. An **invariant Borel class** of countable L -structures is an S_∞ -invariant Borel subset of $\text{Mod}(L)$.

Note that for the purposes of this thesis we always study the orbit equivalence relation, i.e. the isomorphism relation on either $\text{Mod}(L)$ or on the invariant Borel class.

The notion of a Borel reduction allows one to compare complexities of various classification problems.

Definition 4. Let E and F be equivalence relations on the Borel spaces X and Y respectively. E is considered **Borel reducible** to F , denoted $E \leq_B F$, if there exists a Borel function $f : X \rightarrow Y$ such that, for $x, y \in X$, xEy if and only if $f(x)Ff(y)$.

E and F are **bireducible** to each other, denoted $E \sim_B F$, if both $E \leq_B F$ and $F \leq_B E$.

Borel reducibility allows for organization of complexities into a hierarchy. The figure below shows the Borel reductions between the benchmark equivalence relations which are relevant to this thesis.




Figure 1.1: Hierarchy of relevant benchmark equivalence relations

These equivalence relations are increasing in the sense of Borel reducibility from bottom to top wherever there is an edge. We also note that there are many equivalence relations that lie in between those shown which are omitted for the purpose of readability.

At the “bottom” of the Borel hierarchy, we have $\text{id}(\omega)$ which is defined as the equality relation on elements of ω . While this equivalence relation will not be used, this gives one a good base as $\text{id}(\omega)$ is the least complex among equivalence relations with infinitely many classes.

We then see that there exists a Borel reduction from $\text{id}(\omega)$ to $\text{id}(2^\omega)$, or equality on an uncountable Polish space. Note that, for an equivalence relation E on a set X , E is said to be **smooth** if it is the case that E is Borel reducible to $\text{id}(2^\omega)$.

In addition, we see that $\text{id}(\omega)$ is less complex than E_{ω_1} , or isomorphism on codes for countable ordinals.

Increasing in complexity, we arrive at the equivalence relation E_0 which is the immediate successor to $\text{id}(2^\omega)$, among the Borel equivalence relations.

Definition 5. *The equivalence relation E_0 is the relation of eventual equality on 2^ω . That is,*

$$xE_0y \Leftrightarrow \exists m \forall n \geq m x(n) = y(n).$$

At the “top” of our Borel hierarchy, we have Borel complete which is the maximum possible complexity among isomorphism problems for countable structures.

Definition 6. *Given C an invariant Borel class, C is **Borel complete** if and only if, for every invariant Borel class B , B is Borel reducible to C .*

Some commonly referenced examples of Borel complete classes include the class of linear orders [3] and the class of directed graphs [4].

The central aim of this research is to determine the Borel complexities of vertex-transitive structures. We say that a structure A is **vertex-transitive** if, for every element $x, y \in A$, there exists an automorphism φ of A such that $\varphi(x) = y$. Interest in such structures stems from their notable symmetry. In this thesis, we explore the complexities of vertex-transitive graphs, partial orders, linear orders and tournaments whose relationships can be seen below.




Figure 1.2: Structures studied in this thesis

It is important to remark that the class of vertex-transitive structures need not be Borel. To say that the isomorphism relation for such structures is Borel complete, it is sufficient to show that some Borel complete equivalence relation is Borel reducible to it. Formally, we give the following definitions of the structures seen in the figure above.

Definition 7. A **graph** is a pair of sets (V, E) where V is a set of vertices and E is a set of edges formed by pairs of vertices in V . A **directed graph**, or **digraph**, is a graph in which edges are ordered pairs, where edge (u, v) means that there is a directed edge from vertex u to vertex v .

We note that, for our purposes, directed graphs have no self-loops and no bidirectional edges. Further, we see that all other structures can be realized as directed graphs. Narrowing our focus slightly, we find partial orders and tournaments.

Definition 8. A **partial order** is a binary relation \leq on a set X which is:

- i. reflexive: $x \leq x$ for all $x \in X$*
- ii. anti-symmetric: $x \leq y$ and $y \leq x$ implies that $x = y$*
- iii. transitive: $x \leq y$ and $y \leq z$ implies $x \leq z$.*

Definition 9. A *tournament* is a directed graph in which every pair of distinct vertices is connected by a single directed edge.

Finally, tournaments which satisfy the properties of partial orders are linear orders, and vice versa.

Definition 10. A *linear order* is a partial order \leq on a set X which also satisfies the comparability axiom. That is, for every $x, y \in X$ either $x \leq y$ or $y \leq x$.

We begin by extending the results regarding vertex-transitive graphs in Clemens' paper titled Isomorphism of homogeneous structures [1]. In this paper, Clemens showed that the isomorphism relation on countable, connected, vertex-transitive, undirected graphs is Borel complete. Moreover, he suggested that one could also prove this for the directed case. Here, we first cover the simpler directed case before filling in the missing components needed to prove the original result for undirected graphs. A consequence of this leads to the result that vertex-transitive partial orders are Borel complete.

This analysis then leads us to ask about the complexity of a special case of directed graphs: linear orders. We have shown that the class of vertex-transitive linear orderings is, in fact, not Borel complete. Further, we see that there exists an absolutely Δ_2^1 reduction from isomorphism on codes for countable ordinals, denoted E_{ω_1} , to isomorphism on vertex-transitive linear orders.

We finish this paper by broadening our view slightly to consider vertex-transitive tournaments. That is, linear orders are a special type of tournament and so it is natural to see where tournaments lie in terms of Borel complexity. Here we show that isomorphism of vertex-transitive tournaments is properly more complex than E_0 however it remains a question as to whether or not this is Borel complete.

Beyond the results given in this thesis it is natural to ask about other vertex-transitive structures. For example, linear orders are a particular type of lattice, which are partial orders. Since we know that isomorphism of linear orders is Borel complete and isomorphism of partial orders is, in fact, Borel complete it is natural to ask if the class of vertex-transitive lattices is Borel complete as well.

Further, instead of vertex-transitive structures we could consider structures with larger automorphism groups. That is, an n -transitive structure is one in which its automorphism group acts transitively on n -tuples of distinct elements. So, we could instead ask the question: is the isomorphism problem for the class of n -transitive structures Borel complete for a given value n ?

CHAPTER 2

VERTEX-TRANSITIVE GRAPHS AND PARTIAL ORDERS

We begin by considering isomorphism of vertex-transitive graphs. The work shown here is an extension of results from Clemens's paper titled Isomorphism of homogeneous structures [1] in which he first showed that the isomorphism problem for countable, connected, vertex-transitive graphs is Borel complete.

The aim is to break up the proof that the isomorphism problems of countable connected vertex-transitive graphs is Borel complete into several components in order to uncover further results which follow from Clemens's ideas. We will first show that the class of extensional graphs is Borel complete. We then see that there exists a Borel reduction from the isomorphism relation on countable graphs to the isomorphism relation on countable, connected vertex-transitive graphs.

The original proof by Clemens focuses on the case where these graphs are undirected with a comment that we could instead consider the directed case given an additional edge restriction. Here, we first look at the directed case in which we form the directed Cayley graph of H , a group generated by the vertices of a countable graph, as directed Cayley graphs are always vertex-transitive.

Definition 11. A **connected graph** is a graph in which there exists a path between every pair of vertices. A directed graph is **weakly-connected** if replacing all directed edges with undirected edges results in a connected graph.

Theorem 12 (Extracted from [1]). *There exists a Borel reduction from countable graphs to countable, weakly-connected, vertex-transitive, directed graphs.*

Proof. Let $\langle v_i \rangle_{i \in \mathbb{N}}$ enumerate the vertices of a countable graph G . Define H to be the group generated freely by the the vertices of G with the stipulation that adjacent vertices commute. That is, let \mathbb{F}_ω be the free group on generators g_i and N be the normal subgroup of \mathbb{F}_ω generated by $\{g_i g_j g_i^{-1} g_j^{-1} : v_i \text{ adj } v_j \text{ in } G\}$. Then, define $H = \mathbb{F}_\omega / N$ and form Γ , the directed Cayley graph of H with generators $\langle g_i \rangle_{i \in \mathbb{N}}$. The vertices of Γ are left cosets of N in \mathbb{F}_ω and two vertices $w_1 N$ and $w_2 N$ are adjacent in Γ if $g_i w_1 N = w_2 N$ with a directed edge from $w_1 N$ to $w_2 N$. Note that each of the generators g_i are in distinct cosets.

In order to produce a code for this structure, we begin by fixing an enumeration $\langle w_i \rangle_{i \in \mathbb{N}}$ of words in \mathbb{F}_ω . For each coset of N we pick a representative for that coset by picking the least i such that w_i is in the given coset. This w_i is the coset representative. Then, enumerate the chosen representatives and define the binary relation on \mathbb{N} encoding Γ according to whether corresponding cosets are adjacent in Γ .

We now aim to show that the map from G to our code Γ is the desired reduction from the isomorphism relation on countable graphs G to the isomorphism relation on vertex-transitive, countable, weakly-connected, directed graphs. To see that Γ is vertex transitive let $w_1 N$ and $w_2 N$ be vertices of Γ and define the map

$$\varphi(wN) = wNw_1^{-1}w_2 = ww_1^{-1}w_2N,$$

an automorphism of Γ sending w_1N to w_2N .

Now, suppose we have two isomorphic graphs G_1 and G_2 with f an isomorphism between them. Let N_1 and N_2 be the normal subgroups in the construction of Γ_1 and Γ_2 respectively and let

$$\varphi(wN_1) = \tilde{w}N_2$$

where, for $w = g_{i_n}^{\sigma_n} \dots g_{i_0}^{\sigma_0}$, $\tilde{w} = g_{f(i_n)}^{\sigma_n} \dots g_{f(i_0)}^{\sigma_0}$. Well, f induces a partial map φ from their respective Cayley graphs Γ_1 to Γ_2 , such that $\varphi(g_iN_1) = g_{f(i)}N_2$ which acts on the cosets of the g_i . We then extend this partial map to an isomorphism on the graphs as a whole. Certainly, φ is a bijection. To see that it is well-defined, note that the map taking w to \tilde{w} is an automorphism of \mathbb{F}_w which sends N_1 to N_2 and so $w_1w_2^{-1} \in N_1$ if and only if $\tilde{w}_1\tilde{w}_2^{-1} \in N_2$.

Now, suppose that $w_1N_1 \rightarrow w_2N_1$ in Γ_1 or, in other words, $g_kw_1N_1 = w_2N_1$. Then φ gives that $(\widetilde{g_kw_1})N_2 = \tilde{w}_2N_2$. Notice that $\widetilde{g_kw_1} = g_{f(k)}\tilde{w}_1$ and so $g_{f(k)}\varphi(w_1N_1) = \varphi(w_2N_1)$. The argument in the other direction is the same and so $w_1N_1 \rightarrow w_2N_1$ in Γ_1 if and only if $\varphi(w_1N_1) \rightarrow \varphi(w_2N_1)$ in Γ_2 .

Finally, suppose that Γ_1 and Γ_2 are isomorphic. We want to show that $G_1 \cong G_2$ which can be achieved by seeing that one can recover G (up to isomorphism) from the isomorphism class of Γ . Recalling that Γ is vertex-transitive, without loss of generality we can fix a vertex in Γ corresponding to N and identify its adjacent vertices, g_kN . Enumerate these adjacent vertices as $\langle u_i \rangle_{i \in \mathbb{N}}$. Let R be the binary relation on this set such that, for $u_i \neq u_j$,

$$u_i R u_j \Leftrightarrow \exists b \neq N \text{ such that } u_i = g_i N \rightarrow b \text{ and } u_j = g_j N \rightarrow b.$$

The aim is to show that $u_i R u_j$ if and only if there exist $p, q \in \mathbb{N}$ with $u_i = g_p N$ and $u_j = g_q N$ such that $v_p \text{ adj } v_q$ in G . Well, if such a pair p, q exists, then g_p and g_q commute in H . Then u_i and u_j are opposite corners of the square including N and $g_q g_p N = g_p g_q N$. Thus $u_i R u_j$. Suppose now that $u_i R u_j$, with $u_i \neq u_j$. Let $u_i = g_p N$ and $u_j = g_q N$ with N and $b = g_n g_q N = g_m g_p N$ as the other two vertices of the square.




Figure 2.1: Depiction of $u_i R u_j$ with $u_i = g_p N$ and $u_j = g_q N$

Thus it must be that

$$g_p^{-1} g_m^{-1} g_n g_q \in N. \quad (2.1)$$

Recalling that words in N must have the sum of the exponents of each generator equal to 0, it must be that $p = q$ and $m = n$ or $p = n$ and $m = q$. As $u_i \neq u_j$ then $g_p N \neq g_q N$ and so it cannot be the case that $p = q$ and $m = n$. Therefore, $g_p = g_n$ and $g_m = g_q$ and so

$$g_p^{-1} g_q^{-1} g_p g_q \in N$$

which means that $v_p \text{ adj } v_q \in G$. Thus we can recover from Γ an isomorphic copy of G by taking the out-neighbors of N as the vertices and R as the edge relation and so the desired Borel reduction exists. \square

With this, we have that the isomorphism relation on countable, connected, directed vertex-transitive graphs is Borel complete. The proof of the undirected case is very similar and so we exclude the redundancies. We also point out the additional cases required by not knowing whether the powers of the g_i in equation (2.1) are 1 or -1 . Additionally, this proof requires that the graph G be extensional.

Definition 13. An *extensional graph* is one in which, for any vertices $v_1 \neq v_2$, there exists a third vertex v_3 such that $v_1 \text{ adj } v_3$ and not $v_2 \text{ adj } v_3$.

Fortunately, the following Lemma due to Mekler [8] tells us that the isomorphism relation on extensional graphs is also Borel complete. To prove this, note that the isomorphism relation of countable \mathcal{L}_0 -structures is Borel complete [3], where \mathcal{L}_0 is the language containing a single binary relation symbol. We then show that there exists a Borel reduction from the isomorphism relation on \mathcal{L}_0 -structures to extensional graphs which gives the desired conclusion.

Lemma 14 (Mekler). *There exists a Borel reduction from the isomorphism relation on \mathcal{L}_0 -structures to the isomorphism relation on extensional graphs. Thus, the class of extensional graphs is Borel complete.*

Proof. Let A be an \mathcal{L}_0 -structure. First, build $G'(A)$ such that its vertices are elements of A and there are two nodes adjacent to each. If A satisfies $a_1 R a_2$, introduce two new vertices $(a_{1,2})_1$ and $(a_{1,2})_2$ such that $a_1 \text{ adj } (a_{1,2})_1$, $(a_{1,2})_1 \text{ adj } (a_{1,2})_2$, and $(a_{1,2})_2 \text{ adj } a_2$. Further, insert three new vertices adjacent to $(a_{1,2})_1$ and four new vertices adjacent

to $(a_{1,2})_2$. The graph of M constructed up to this point can be seen in the figure below.




Figure 2.2: Graph of M as constructed by Mekler, [8]

Finally, insert three new vertices adjacent to one another and to each of the previous vertices of $G'(A)$. Once $G'(A)$ is constructed, let each element of $G'(A)$ be a vertex of $G(A)$. $G(A)$ is constructed by inserting a new vertex in the middle of each edge. Let $\langle v_i \rangle_{i \in \mathbb{N}}$ enumerate the vertices of $G(A)$. We note that $G(A)$ is extensional.

In order to show that this Borel map is, in fact, a Borel reduction we must show that $A_1 \cong A_2$ if and only if $G(A_1) \cong G(A_2)$. If $A_1 \cong A_2$, then there exists an isomorphism $\varphi : A_1 \rightarrow A_2$ such that if $a_1 R a_2$ in A_1 then $\varphi(a_1) R \varphi(a_2)$ in A_2 . As $a_1 R a_2$, we obtain the resulting encoding $G(A_1)$. Applying φ to the vertices representing a_1 and a_2 , we achieve an isomorphism φ' sending vertices in $G(A_1)$ to vertices in $G(A_2)$ and so $G(A_1) \cong G(A_2)$.

Suppose now that $G(A_1) \cong G(A_2)$. That is, there exists an isomorphism $\varphi' : G(A_1) \rightarrow G(A_2)$ sending structures representing adjacent vertices in A_1 to structures representing adjacent vertices in A_2 . Consider any pairs of elements $a_1 \text{ adj } a_2$ in A_1 and $a_3 \text{ adj } a_4$ in A_2 . Then there exists an encoding in $G(A_1)$ of $a_1 \text{ adj } a_2$ and similarly for $G(A_2)$.

To see that we can recover the two vertices representing elements in A_1 , one can locate all vertices which are a distance of two from vertices of degree four in $G(A)$. From these, one can identify v_1 and v_2 representing a_1 and a_2 respectively as follows: v_1 and v_2 have five vertices in between them, where v_1 is a distance of two from a vertex of degree eight and v_2 is a distance of two from a vertex of degree nine. Thus, by our encoding, we can recover that $a_1 \text{ adj } a_2$.

As these graphs are isomorphic, we get that there exists an isomorphism φ such that $v_1 R v_2$ with respect to the encoding if and only if $\varphi(v_1) R \varphi(v_2)$ or, equivalently, $v_3 R v_4$. Thus, we arrive at $a_1 \text{ adj } a_2$ if, and only if $a_3 \text{ adj } a_4$ and so $A_1 \cong A_2$ as expected. \square

Theorem 15 (Clemens). *There exists a Borel reduction from extensional graphs G to countable, connected, undirected, vertex-transitive graphs.*

Proof. The proof of this result very closely parallels the proof of Theorem 12, except we form Γ , the undirected Cayley graph of H , where two vertices $w_1 N$ and $w_2 N$ are adjacent in Γ if there exists a generator g_i such that $g_i w_1 N = w_2 N$ or $g_i w_2 N = w_1 N$. We then continue to show that the map $G \mapsto \Gamma$ is the desired reduction. Showing that $G_1 \cong G_2$ implies $\Gamma_1 \cong \Gamma_2$ is the same as in the proof of Theorem 12.

In order to see that $\Gamma_1 \cong \Gamma_2$ gives $G_1 \cong G_2$ requires showing that G can be recovered from Γ . This becomes slightly more involved than in the directed case. The argument begins the same; however, instead of the definition of the relation R given above, for two distinct elements u_i and u_j in Γ we define R as

$$u_i R u_j \Leftrightarrow \exists a \exists b, a \neq b, \text{ such that } u_i \text{ and } u_j \text{ are each adjacent to both } a \text{ and } b$$

We then aim to show that $u_i R u_j$ if and only if there exist $k_1, k_2 \in \mathbb{N}$ and $\sigma_1, \sigma_2 \in \{-1, 1\}$ with $u_i = g_{k_1}^{\sigma_1} N$ and $u_j = g_{k_2}^{\sigma_2} N$ such that $v_{k_1} \text{ adj } v_{k_2}$ in G .

Well, if such a pair k_1, k_2 exists then g_{k_1} and g_{k_2} commute in H . Thus u_i and u_j are at opposite corners of a square shared with N and $g_{k_1}^{\sigma_1} g_{k_2}^{\sigma_2} N = g_{k_2}^{\sigma_2} g_{k_1}^{\sigma_1} N$ and so $u_i R u_j$.




Figure 2.3: Resulting square if there exist $k_1, k_2 \in \mathbb{N}$ with $v_{k_1} \text{ adj } v_{k_2} \in G$

Now, suppose that $u_i R u_j$ with $u_i = g_{k_1}^{\sigma_1} N$ and $u_j = g_{k_2}^{\sigma_2} N$ and a, b the other two vertices of the square. Thus there must exist generators $g_{n_1}, g_{n_2}, g_{m_1}, g_{m_2}$ and $\tau_1, \tau_2, \rho_1, \rho_2 \in \{-1, 1\}$ such that

$$a = g_{n_1}^{\tau_1} g_{k_1}^{\sigma_1} N = g_{n_2}^{\tau_2} g_{k_2}^{\sigma_2} N \text{ and } b = g_{m_1}^{\rho_1} g_{k_1}^{\sigma_1} N = g_{m_2}^{\rho_2} g_{k_2}^{\sigma_2} N.$$

So we have

$$g_{k_1}^{-\sigma_1} g_{n_1}^{-\tau_1} g_{n_2}^{\tau_2} g_{k_2}^{\sigma_2} \in N \text{ and } g_{k_1}^{-\sigma_1} g_{m_1}^{-\rho_1} g_{m_2}^{\rho_2} g_{k_2}^{\sigma_2} \in N.$$

As words in N must have the sum of the exponents of each generator equal to 0, then it must be that $k_1 = k_2$, $k_1 = n_1$, or $k_1 = n_2$. If $k_1 = k_2$ then it must be that $-\sigma_1 = \sigma_2$. Otherwise, $u_i = g_{k_1}^{\sigma_1} N$ and $u_j = g_{k_2}^{\sigma_2} N$ would be the same. Further, this would require that $n_1 = n_2 = k_1 = k_2$ and so $\sigma_2 - \sigma_1 + \tau_2 - \tau_1 = 0$. Well, then we would need $-\sigma_1 = \tau_1$ and so $a = g_{k_1}^{-\sigma_1} g_{k_1}^{\sigma_1} N = N$. In the case that $k_1 \neq k_2$ and $k_1 = n_1$, we get that $-\sigma_1 = \tau_1$ and so, once again, $a = g_{k_1}^{-\sigma_1} g_{k_1}^{\sigma_1} N = N$. Finally, for the case where $k_1 \neq k_2$ and $k_1 = n_2$ it also must be the case that $k_2 = n_1$, $\sigma_1 = \tau_2$, and $\sigma_2 = \tau_1$. Well, then $g_{k_1}^{-\sigma_1} g_{k_2}^{-\sigma_2} g_{k_1}^{\sigma_1} g_{k_2}^{\sigma_2} \in N$ and so $v_{k_1} \text{ adj } v_{k_2}$ in G .

Repeating the same argument for b , we see that either $b = N$ or $v_{k_1} \text{ adj } v_{k_2}$ in G . Recalling that $a \neq b$ it must be the case that $v_{k_1} \text{ adj } v_{k_2}$ in G . Thus we can identify pairs of elements $\{u_i, u_j\}$, so that they are R -related to all of the same elements. That is, we have identified pairs $\{g_k N, g_k^{-1} N\}$. By extensionality, this will not identify any other pairs as, given g_i and g_j , $i \neq j$, there exists a g_k which commutes with g_i but not with g_j and vice versa.

Therefore, we form the graph with these pairs as its vertices where two vertices are set adjacent if each element in the first pair is R -related to each element in the second pair. The relation R then gives that this graph is isomorphic to G . \square

Therefore, the isomorphism relation on countable, connected, vertex-transitive graphs is Borel complete. A further consequence is that vertex-transitive partial orders are also Borel complete. This can be seen by taking the transitive closure of the directed graph from Theorem 12 which will result in a partial order, as we will do below. First, we recall the definition of a partial order:

Definition 16. A *partial order* is a relation which is reflexive, antisymmetric, and transitive.

Definition 17. The *transitive closure* of a directed graph G , denoted $C(G)$, is a graph which contains an edge $\{u, v\}$ whenever there is a directed path from u to v , for $u, v \in G$ as well as a path of length 0 from u to u .

It is important to note that the transitive closure of a directed graph does not always yield a partial order. For this to be the case, it must be that the directed graph has no cycles. For the following statement, we remind the reader that, in this paper, directed graphs have no cycles of length one or two.

Proposition 18. *If G a directed graph, then the transitive closure $C(G)$, is a partial order if, and only if, G has no directed cycles.*

Proof. For the forward direction, suppose $C(G)$ satisfies the requirements of a partial order. Further, suppose towards a contradiction that G contains a cycle such that $x_1 \leq x_2 \leq \dots \leq x_n \leq x_1$ for x_1, x_2, \dots, x_n vertices of $G(A)$. However, as $C(G)$ is antisymmetric, $x_1 \leq x_i$ and $x_i \leq x_1$ gives $x_1 = x_i$ for $1 \leq i \leq n$. Therefore, it cannot be the case that G has a cycle.

Now, suppose there exist no cycles in G . To show that $C(G)$ satisfies the requirements of a partial order, we first note that the transitive closure is, in fact, transitive. Further, we assumed the relation on A was reflexive and so $C(G)$ remains transitive, as this does not affect single vertices but the relationship between two vertices. And finally, as there are no cycles in G it must be that there are no cycles in $C(G)$ and so we achieve antisymmetry. Thus the transitive closure of G admits a partial order. \square

Corollary 19. *There exists a Borel reduction from the isomorphism relation on extensional graphs G to the isomorphism relation on countable vertex-transitive partial orders.*

Proof. Recalling Proposition 18, note that the directed graph Γ of Theorem 12 has no directed cycles and so the transitive closure, $C(\Gamma)$, is a partial order. To see that Γ has no directed cycles, note that Γ is the directed Cayley graph of H in which, for all words $w = w_1 \dots w_k$ having generators g_1, \dots, g_k , if $w = 1$ in H then the sum of all exponents of each of the corresponding g_i must equal 0. Therefore, the sum is not positive and so there cannot be a directed cycle.

Therefore, given Γ one can produce its transitive closure, $C(\Gamma)$, which is a partial order. The aim is to show that $\Gamma_1 \cong \Gamma_2$ if and only if $C(\Gamma_1) \cong C(\Gamma_2)$. If $\Gamma_1 \cong \Gamma_2$ then we certainly have that $C(\Gamma_1) \cong C(\Gamma_2)$ as transitive closure is isomorphism invariant. So, now suppose that $C(\Gamma_1) \cong C(\Gamma_2)$. Given $C(\Gamma)$ we want to show that we can recover the original $b \in \Gamma$ such that $N \rightarrow b$ in Γ from any point $p \in C(\Gamma)$.

The claim is that the out-neighbors of N are all of the $b \in \Gamma$ such that $N \rightarrow b$ in $C(\Gamma)$ and there does not exist a directed path of length greater than one from N to b in $C(\Gamma)$. This is the case as, if there did exist such a path in $C(\Gamma)$ then we would have

$$b = g_j N \text{ and } b = g_{i_1} \dots g_{i_n} N$$

which would mean that $g_{i_1} \dots g_{i_n} g_j^{-1} = 1$ in H which contradicts that words in N must have the sum of the exponents of all generators equal to 0. Thus we have recovered Γ from $C(\Gamma)$ and so there exists a Borel reduction from extensional graphs to countable partial orders. \square

CHAPTER 3

VERTEX-TRANSITIVE LINEAR ORDERINGS

After classifying directed graphs in the previous chapter it is natural to ask about the complexity of special cases of directed graphs such as linear orders. The main theorem of this chapter isolates the complexity of vertex-transitive linear orders, in particular. Before presenting this result, it is necessary to discuss the condensation of linear orders. Condensation will be used in the main theorem in order to classify vertex-transitive linear orders by countable ordinals (refer to [9] for more details).

First, let us recall that a linear order is a partial order \leq on a set X which also satisfies comparability: for every $x, y \in X$, either $x \leq y$ or $y \leq x$.

Definition 20. *Let L be a linear ordering and let L' be a collection of non-empty intervals of L which partitions L , ordered by*

$$I_1 \ll I_2 \text{ if, for all } x_1 \in I_1, x_2 \in I_2, x_1 < x_2.$$

*L' is called a **condensation** of L .*

Intuitively, we think of the condensation as whole intervals of L being condensed to single points in L' . It is also important to note that the condensation of a vertex-transitive linear ordering remains vertex-transitive. That is, if L is vertex-transitive then there exists an automorphism φ such that $\varphi(x) = y$ for every pair $x, y \in L$

and this automorphism extends to $C[L]$. For our purposes, we will make use of finite condensation maps, where discrete intervals are condensed.

Definition 21. *The **finite condensation map** of L is denoted*

$$c_F(x) = \{y \mid [x, y] \text{ or } [y, x] \text{ is finite}\}$$

for $x \in L$. Let $c_F[L]$ denote the condensation of L determined by the intervals $c_F(x)$ for all $x \in L$.

Note that $c_F(x)$ can be finite or, otherwise, has order type ω , ω^* or \mathbb{Z} . We call $c_F(x)$ the equivalence class of x . This condensation map will be used implicitly throughout this chapter. Considering more than a single condensation leads to the following iterative definitions.

Definition 22. *We define an **iterated condensation map** with $c^1 = c$ as*

$$(i) \ c^{\alpha+1}(x) = \{y \mid c(c^\alpha(x)) = c(c^\alpha(y))\}$$

(ii) *Assume that λ is a limit ordinal and that for each $\beta < \lambda$ we have defined for each linear ordering A the condensation map c^β . Then we define the condensation map c^λ by*

$$c^\lambda(x) = \bigcup \{c^\beta(x) \mid \beta < \lambda\}$$

With the above definition of iterated condensations we note that, given a linear order L , the condensation map stabilizes after a bounded number of iterations. The following proposition shows that this is, in fact, the case.

Proposition 23. *Let L be a linear ordering of cardinality κ . Then there exists an ordinal $\alpha < \kappa^+$ such that $c^\beta(x) = c^\alpha(x)$ for all $x \in L$ and for all $\beta \geq \alpha$.*

Proof. If $c^{\alpha+1}(x) = c^\alpha(x)$ for every $x \in L$, then $c^\beta(x) = c^\alpha(x)$ for every $\beta \geq \alpha$. Thus, for some set $S = \{\alpha \mid c^{\alpha+1}(x) \neq c^\alpha(x) \text{ for some } x \in L\}$, S is an initial segment of ordinals.

Further, for each $\alpha \in S$, there exists a pair $\{x, y\}$ of elements of L such that $c^\alpha(x) \neq c^\alpha(y)$ but $c^{\alpha+1}(x) = c^{\alpha+1}(y)$. So S is an initial segment of the ordinals which is in a one-to-one correspondence with a subset of $L \times L$. Thus S has cardinality at most κ .

Therefore, α has cardinality κ and so $\kappa \leq \alpha \leq \kappa^+$ [9]. Thus we have an $\alpha < \kappa^+$ such that $c^\beta(x) = c^\alpha(x)$ for every $x \in L$ and for every $\beta \geq \alpha$. \square

Recalling that our aim is to understand vertex-transitive linear orders, we will show that there are two such classes of vertex-transitive linear orders, namely powers of \mathbb{Z} and \mathbb{Q} copies of powers of \mathbb{Z} . This leads us to the following definition:

Definition 24. *Given an ordinal β , $\mathbb{Z}_0(\beta)$ consists of all β -sequences of elements of \mathbb{Z} which have only finitely many non-zero entries. That is*

$$\mathbb{Z}_0(\beta) = \{s : \beta \rightarrow \mathbb{Z} \mid \{\alpha : s(\alpha) \neq 0\} \text{ is finite}\},$$

ordered by $s \ll t$ if $s(\mu) < t(\mu)$ where μ is the largest ordinal $\gamma < \beta$ for which $s(\gamma) \neq t(\gamma)$.

For a more intuitive definition of powers of \mathbb{Z} , one can recursively construct a definition of $\mathbb{Z}_0(\beta)$, which we will call \mathbb{Z}^β , as follows:

Definition 25. (i) $\mathbb{Z}^0 = 1$

$$(ii) \mathbb{Z}^{\beta+1} = \mathbb{Z}^\beta \cdot \omega^* + \mathbb{Z}^\beta + \mathbb{Z}^\beta \cdot \omega = \mathbb{Z}^\beta \cdot \mathbb{Z}$$

$$(iii) \mathbb{Z}^\lambda = (\sum \{\mathbb{Z}^\gamma \cdot \omega \mid \gamma < \lambda\})^* + 1 + \sum \{\mathbb{Z}^\gamma \cdot \omega \mid \gamma < \lambda\} \text{ for limit ordinals } \lambda.$$

Note that we define the **sum** of arbitrarily many linear orderings as follows. Let $\langle I, R \rangle$ be a linear ordering and, for each $i \in I$, let $\langle A_i, S_i \rangle$ be a linear ordering. Then $\sum_{i \in I} A_i$ is defined to be the linear ordering $\langle C, T \rangle$ with $C = \bigcup_{i \in I} A_i$ and, for any two $c_1, c_2 \in C$

$$c_1 <_T c_2 \text{ if } (c_1 \in A_i \text{ and } c_2 \in A_j \text{ and } i <_R j) \text{ or} \\ (c_1 \in A_i \text{ and } c_2 \in A_i \text{ and } c_1 <_{S_i} c_2 \text{ for some } i \in I).$$

Further, we define the **product** of two linear orders, $\langle I, R \rangle$ and $\langle A, S \rangle$ as

$$A \cdot I = \sum_{i \in I} A$$

where addition is as defined above.

The following proposition tells us that the previous definitions of powers of \mathbb{Z} are, in fact, equivalent.

Proposition 26. For any ordinal β , $\mathbb{Z}_0(\beta) \cong \mathbb{Z}^\beta$.

Proof. This is proved by induction on β . First, consider the case $\mathbb{Z}_0(0) \cong \mathbb{Z}^0$. By definition $\mathbb{Z}^0 = 1$ and so now $\mathbb{Z}_0(0) = \{s : 0 \rightarrow \mathbb{Z} \mid \exists N \forall n \geq N s(n) = 0\} \cong \{\emptyset\} \cong 1$. Thus $\mathbb{Z}^0 = 1 \cong \mathbb{Z}_0(0)$.

Now, suppose we have that $\mathbb{Z}_0(\beta) = \mathbb{Z}^\beta$ and let us show that the successor case holds, i.e. $\mathbb{Z}_0(\beta + 1) \cong \mathbb{Z}^{\beta+1}$ below:

$$\begin{aligned}
\mathbb{Z}^{\beta+1} &= \mathbb{Z}^\beta \cdot \mathbb{Z} \\
&\cong \mathbb{Z}_0(\beta) \cdot \mathbb{Z} \text{ (by assumption)} \\
&\cong \{(s : (\beta) \rightarrow \mathbb{Z}, t \in \mathbb{Z}) \mid \exists N \forall n \geq N s(n) = 0\} \\
&= \mathbb{Z}_0(\beta + 1).
\end{aligned}$$

Finally, for a limit ordinal λ , we claim that $\mathbb{Z}_0(\lambda) = \{s : \lambda \rightarrow \mathbb{Z} \mid \{\alpha : s(\alpha) \neq 0\} \text{ is finite}\}$ and $\mathbb{Z}^\lambda = (\sum \{\mathbb{Z}^\gamma \cdot \omega \mid \gamma < \lambda\})^* + 1 + \sum \{\mathbb{Z}^\gamma \cdot \omega \mid \gamma < \lambda\}$ are isomorphic.

To see this, let $\circlearrowleft \in \mathbb{Z}_0(\lambda)$ denote the sequence $s(\alpha) = 0$ for all α . This \circlearrowleft corresponds to the “1” in the definition of \mathbb{Z}^λ . Next, the $s \in \mathbb{Z}_0(\lambda)$ such that $s(0) > 0$ and $s(\alpha) = 0$ for all $\alpha > 0$ are a copy of $\mathbb{Z}^0 \cdot \omega = \omega$.

In general, the elements $s \in \mathbb{Z}_0(\lambda)$ with $s(\gamma) > 0$ and $s(\alpha) = 0$ for all $\alpha > \gamma$ are a copy of $\mathbb{Z}^\gamma \cdot \omega$. Thus, we account for the sum $\sum \{\mathbb{Z}^\gamma \cdot \omega \mid \gamma < \lambda\}$ in the definition of \mathbb{Z}^λ .

There is an analogous correspondence between sequences in $\mathbb{Z}_0(\lambda)$ and the sum $(\sum \{\mathbb{Z}^\gamma \cdot \omega \mid \gamma < \lambda\})^*$. Thus, $\mathbb{Z}_0(\lambda) \cong \mathbb{Z}^\lambda$ for λ a limit ordinal.

Therefore we conclude that, for any ordinal β , $\mathbb{Z}_0(\beta) \cong \mathbb{Z}^\beta$. □

To understand how condensations of power of \mathbb{Z} behave, let us first look at a single condensation of $\mathbb{Z}^{\beta+1}$ where $\beta + 1$ is finite.

Proposition 27. *A single condensation of $\mathbb{Z}^{\beta+1}$, for $\beta + 1 < \omega$, reduces to \mathbb{Z}^β .*

Proof. By definition of finite condensation, we see that

$$c[\mathbb{Z}^{\beta+1}] = \{\{(a, b) | b \in \mathbb{Z}\} | a \in \mathbb{Z}^\beta\} \cong \mathbb{Z}^\beta$$

and so a single condensation of $\mathbb{Z}^{\beta+1}$ reduces to \mathbb{Z}^β in the case that $\beta + 1 < \omega$. \square

Moreover, given sequences in \mathbb{Z}^λ for some limit ordinal λ we see that condensations of such sequences behave as described in the following proposition.

Proposition 28. *For $\mathbb{Z}^\lambda = \{s : \lambda \rightarrow \mathbb{Z} \mid \{\alpha : s(\alpha) \neq 0\} \text{ is finite}\}$, given two sequences $s, t \in \mathbb{Z}^\lambda$*

$$(i) \ c(s) = \{t \mid s \upharpoonright_{[1, \lambda]} = t \upharpoonright_{[1, \lambda]}\}$$

$$(ii) \ c^\gamma(s) = \{t \mid s \upharpoonright_{[\gamma, \lambda]} = t \upharpoonright_{[\gamma, \lambda]}\} \text{ for each ordinal } \gamma.$$

Proof. Let $\mathbb{Z}^\lambda = \{s : \lambda \rightarrow \mathbb{Z} \mid \{\alpha : s(\alpha) \neq 0\} \text{ is finite}\}$ and consider two sequences $s, t \in \mathbb{Z}^\lambda$.

- (i) First, suppose $s \upharpoonright_{[1, \lambda]} = t \upharpoonright_{[1, \lambda]}$ and, without loss of generality, let $s \ll t$. Then we must have $s(0) < t(0)$. If there exists a third $r \in \mathbb{Z}^\lambda$ and r differs at a latter point γ then it does from both s and t so either $r \ll s \ll t$ or $s \ll t \ll r$. If $s \ll r \ll t$ then it must be that $s(0) < r(0) < t(0)$ and there are only finitely many r such that this is the case. Therefore $[s, t]$ is finite and so $c(s) = c(t)$.

Now, if we suppose that $s \upharpoonright_{[1, \lambda]} \neq t \upharpoonright_{[1, \lambda]}$ then there exists a last ordinal γ such that $1 \leq \gamma < \lambda$ where $s(\gamma) \neq t(\gamma)$. Without loss of generality, let $s(\gamma) < t(\gamma)$. Let $r_n \in \mathbb{Z}^\lambda$ be defined as $r_n = n \frown s \upharpoonright_{[1, \lambda]}$ for any $n \in \mathbb{Z}$. Then $s \ll r_n \ll t$ for

$n > s(0)$ and there are infinitely many r_n which satisfy this. Therefore, $[s, t]$ is infinite and so $c(s) \neq c(t)$.

Therefore, we conclude that $c(s) = \{t \mid s \upharpoonright_{[1,\lambda)} = t \upharpoonright_{[1,\lambda)}\}$.

(ii) This case follows using a similar argument to (i) and induction on γ .

□

As we alluded to earlier in the chapter, there are two classes of vertex-transitive linear orders. In order to prove this, we require the following definitions about single elements of L .

Definition 29. *Given a linear ordering L ,*

(i) $x \in L$ is **left dense** if x is not the least element in L and if there is no greatest element $y < x$

(ii) $x \in L$ is **right dense** if x is not the greatest element in L and if there is no least element $y > x$

(iii) $x \in L$ is **left discrete** if there exists a greatest $y \in L$ such that $y < x$ or if x is the greatest element of L

(iv) $x \in L$ is **right discrete** if there exists a least $y \in L$ such that $y > x$ or if x is the least element of L .

Note that, given some element $x \in L$, x will be either left/right dense or left/right discrete. This leads us to the following lemma, where we will see that \mathbb{Z}^β and $\mathbb{Z}^\beta \cdot \mathbb{Q}$, for countable ordinals β , (along with the trivial linear order 1) are the only possible classes of vertex-transitive linear orders.

Lemma 30. *If a linear ordering L is vertex-transitive, then*

- (i) *if one point is left or right dense then every point is and, in fact, $L \cong \mathbb{Q}$*
- (ii) *if one point is left or right discrete then every point is, and every equivalence class is a copy of the integers or $L = 1$.*

Proof. By vertex-transitivity, if one point is left or right dense then every point must also be left or right dense, respectively.

- (i) First, let $a < b$. If b is left dense then, given a sequence $\{b_i\}_{i \in \mathbb{N}}$ approaching b from the left, there must eventually be some $n \in \mathbb{N}$ such that $a < b_n < b$. Therefore, L is dense. Similarly, let $b < a$ be right dense. Then, given a sequence $\{b_i\}_{i \in \mathbb{N}}$ approaching b from the right, there must eventually be some $n \in \mathbb{N}$ such that $b < b_n < a$. Once again, L is dense.

As L is a dense linear ordering, then L is isomorphic to \mathbb{Q} , $\mathbb{Q} \cup \{\infty\}$, $\{-\infty\} \cup \mathbb{Q}$, or $\{-\infty\} \cup \mathbb{Q} \cup \{\infty\}$. However, as L is also vertex-transitive there can be no first/last element and so it must be the case that $L \cong \mathbb{Q}$.

- (ii) Similar to above, vertex-transitivity tells us that if one point is left or right discrete then every point must be left or right discrete. Thus L is discrete and so every equivalence class is either a single point, a finite number of points, ω , ω^* , or \mathbb{Z} . As before, the vertex-transitivity of L requires that there exists no first/last element of any class and so we must have that every equivalence class is a copy of the integers in the case that L has more than one element.

□

Following Proposition 23, which states that we reach a condensation fixed-point after a bounded number of iterations, and the discussion of iterated condensations of

vertex-transitive linear orders, it is natural to ask what happens when we reach such a fixed-point.

Proposition 31. *Given a vertex-transitive linear order L , if $c(x) = \{x\}$ for every $x \in L$ then L is isomorphic to either \mathbb{Q} or 1 .*

Proof. Let L be a vertex-transitive linear order and $c(x) = \{x\}$ for every $x \in L$.

There are two cases to consider:

- (i) If one point is left/right dense then, by Lemma 30, $L \cong \mathbb{Q}$.
- (ii) If one point is left/right discrete then Lemma 30 tells us that there are two possibilities. The first is that $L = 1$. Otherwise, every equivalence class of L is a copy of the integers. However, if this is the case then, for any $x \in L$, $c(x) = \{y \mid [x, y] \text{ or } [y, x] \text{ is finite}\} \neq \{x\}$ as every $x \in \mathbb{Z}$ is a finite distance from at least one other $y \in \mathbb{Z}$. Thus we reach a contradiction and so if one point is left/right discrete at $c(x) = \{x\}$ for every $x \in L$ then $L = 1$.

□

By Proposition 23 we are able to conclude that for countable, vertex-transitive linear orders it takes countably many steps to arrive at a condensation fixed point. Further, by Proposition 31 the possible condensation fixed points are either 1 or \mathbb{Q} . In summary, we arrive at the following theorem.

Theorem 32. *If L is a countable vertex-transitive linear order then L is isomorphic to either \mathbb{Z}^β or $\mathbb{Z}^\beta \cdot \mathbb{Q}$ for some $\beta < \omega_1$.*

To conclude this chapter, we find that there exists an absolutely Δ_2^1 reduction from isomorphism of codes for countable ordinals to isomorphism of vertex-transitive linear orders.

Theorem 33. *There exists an absolutely Δ_2^1 reduction from isomorphism on codes for countable ordinals to isomorphism on vertex-transitive linear orders.*

Proof. The aim is to show that there exists a Δ_2^1 map from codes for ordinals α to codes for \mathbb{Z}^α . We can construct such a map by recursion on α . Given a code $<_L$ on ω for L we define, for $n, m \in L \cdot \mathbb{Z}$ where $n = \langle n_0, n_1 \rangle$ and $m = \langle m_0, m_1 \rangle$, $n <_{L \cdot \mathbb{Z}} m$ if and only if $n_0 <_{\mathbb{Z}} m_0$ or $n_0 = m_0$ and $n_1 <_L m_1$. For limit stages, given a code for a limit ordinal λ , together with a λ -sequence of codes for \mathbb{Z}^α , $\alpha < \lambda$, we can produce a code for $\mathbb{Z}^\lambda = (\sum \{\mathbb{Z}^\gamma \cdot \omega \mid \gamma < \lambda\})^* + 1 + \sum \{\mathbb{Z}^\gamma \cdot \omega \mid \gamma < \lambda\}$ in a similar manner.

It is not difficult to check that each step of this recursive construction is Borel. Further, it is well known that this implies we can construct a code for \mathbb{Z}^β in an absolutely Δ_2^1 fashion. For example, an infinite time turing machine (ITTM) can easily be programmed to carry out the recursive construction, and ITTM-computable mappings are always absolutely Δ_2^1 (for the definition of ITTM and the statement of this fact, see [5]). □

CHAPTER 4

VERTEX-TRANSITIVE TOURNAMENTS

We now shift focus to the classification problem for vertex-transitive tournaments. As tournaments are a broader class than linear orders, but still a subset of the class of directed graphs it makes sense to consider the complexity of these structures as well. While the exact complexity of vertex-transitive tournaments is yet to be determined, the main result in this chapter states that isomorphism of vertex-transitive tournaments is properly more complex than E_0 , or eventual equality on 2^ω . A remaining question is to determine whether this is, in fact, Borel complete.

Definition 34. A *tournament* is a directed graph in which every pair of vertices is connected by an edge.

Recall that a tournament is vertex-transitive if its automorphism groups acts transitively on the set of vertices. We denote the isomorphism relation over vertex-transitive tournaments as \cong_{VTT} .

Further, we note the following two relations which will be necessary in the proof of the main theorem. First, eventual equality on 2^ω is denoted E_0 . More formally, we say

$$xE_0y \Leftrightarrow \exists m \forall n \geq m x(n) = y(n)$$

We will find that, instead of working with E_0 directly, we will need to work with a subset of the domain of $E_{\mathbb{Z}}$, or the shift equivalence on $2^{\mathbb{Z}}$. That is,

$$xE_{\mathbb{Z}}y \Leftrightarrow \exists m \forall n x(n+m) = y(n)$$

In order to show that isomorphism of vertex-transitive tournaments is more complex than E_0 we will see that the necessary reduction only works on a subset of the domain of $E_{\mathbb{Z}}$. However, this makes it necessary to check that the restriction of $E_{\mathbb{Z}}$ to this subset remains as complex as E_0 which leads us to the following proposition.

Proposition 35. *E_0 on 2^{ω} is Borel bireducible to $E_{\mathbb{Z}} \upharpoonright_{\mathcal{C}}$ on $2^{\mathbb{Z}}$ for any comeager set \mathcal{C} .*

Proof. First, we claim that $E_{\mathbb{Z}}$ has a dense orbit. To see this, note that $2^{\mathbb{Z}}$ has the topology with basic open sets $V_t = \{x \in 2^{\mathbb{Z}} \mid t \subset x\}$. Recall that $2^{<\omega}$ is countable and let $\{s_i\}_{i \in \omega}$ enumerate $2^{<\omega}$. Further, we define $x = \emptyset \frown s_1 \frown s_2 \frown \dots \frown s_i \frown \dots$ where \emptyset denotes the sequence of all 0 on ω^* . This x is certainly in $2^{\mathbb{Z}}$ and contains an instance of every finite binary sequence. That is, given $s \in 2^{<\omega}$, there exists some i such that $s = s_i = x(j) \upharpoonright_{|s|}$ where $|s|$ denotes the length of the sequence s_i .

Further, given $s \in 2^{<\omega}$ as before, we know that $s \subset z \cdot x$ as $s = x(j-z) \upharpoonright_{|s|}$. Therefore, shifts of x also contain an instance of every finite binary sequence.

The aim is to show that, given $[x]_{\mathbb{Z}} = \{z \cdot x \mid z \in \mathbb{Z}\}$ where $z \cdot x = x(n+z)$ for all $n \in \mathbb{Z}$, every V_t contains an element of $[x]_{\mathbb{Z}}$.

That is, given any V_t , we know that $t \subset z \cdot x$ for some $z \in \mathbb{Z}$ and so $t \in [x]_{\mathbb{Z}}$. Therefore $t \in V_t \cap [x]_{\mathbb{Z}}$ and so there exists a dense orbit. As this is the case, we know that $E_{\mathbb{Z}}$ is generically ergodic, meaning that every $E_{\mathbb{Z}}$ -invariant Borel subset of $2^{\mathbb{Z}}$ is either meager or comeager (Proposition 6.1.9, [4]).

As $E_{\mathbb{Z}}$ is generically ergodic, we also know that $E_{\mathbb{Z}} \upharpoonright_{\mathcal{C}}$ is generically ergodic (Proposition 6.1.9, [4]). As this is the case and all orbits of \mathbb{Z} are countable, hence meager, we find that $E_{\mathbb{Z}} \upharpoonright_{\mathcal{C}}$ is not smooth (Proposition 6.1.10, [4]). As $E_{\mathbb{Z}} \upharpoonright_{\mathcal{C}}$ is not smooth, one can conclude that E_0 is Borel reducible to $E_{\mathbb{Z}} \upharpoonright_{\mathcal{C}}$ (Proposition 6.3.1, [6]). On the other hand, we know that any \mathbb{Z} -orbit relation is Borel reducible to E_0 [2] and therefore $E_{\mathbb{Z}} \upharpoonright_{\mathcal{C}}$ is Borel bireducible with E_0 . \square

Thus we arrive at the main result for vertex-transitive tournaments. That is, the isomorphism relation for vertex-transitive tournaments is more complex than E_0 .

Theorem 36. *There exists a Borel reduction from E_0 to isomorphism of vertex-transitive tournaments.*

Proof. Given $x \in 2^{\mathbb{Z}}$, we construct a tournament, T_x , with vertices in $\mathbb{Z} \times \mathbb{Z}$ as follows.


Given two vertices at positions (m, n) and (m', n') where $m, n, m', n' \in \mathbb{Z}$, we define

$$(m, n) \rightarrow (m', n') \Leftrightarrow \begin{cases} m = m' \text{ and } n > n' \text{ or} \\ m' = m + 1 \text{ and } x(n' - n) = 1 \text{ or} \\ |m - m'| \geq 2 \text{ and } m < m' \end{cases}$$

where, otherwise, there is an edge $(m', n') \rightarrow (m, n)$. An example of this can be seen in the figure below, where we focus on a pair of columns which are “one apart”.

Note that x is bi-infinite, but for illustration purposes we highlight five consecutive terms in x and the five corresponding vertices in relation to vertex $(0, 0)$.

It is important to note that these T_x are, in fact, vertex-transitive as the automorphism carrying the vertex at position $(0, 0)$ to position (m, n) is given by the map $(a, b) \mapsto (a + m, b + n)$ preserves the definition of edge direction given above.



n	...	-2	-1	0	1	2	...
$x(n)$...	1	1	0	0	1	...

Figure 4.1: Directed edges from vertex $(0,0)$ to vertices one column to the right.

Now that we have developed the construction of T_x given x , the aim is to show that there exists a comeager subset $\mathcal{C} \subseteq 2^{\mathbb{Z}}$, which will be defined later, such that, for all sequences $x, x' \in \mathcal{C}$,

$$xE_{\mathbb{Z}}x' \Leftrightarrow T_x \cong T_{x'}.$$

To begin, suppose that $xE_{\mathbb{Z}}x'$. That is, there exists some $k \in \mathbb{Z}$ such that, for every $n \in \mathbb{Z}$, $x(n) = x'(n+k)$. Now, construct two vertex-transitive tournaments, T_x and $T_{x'}$, from x and x' respectively as described above. In order to see that T_x and $T_{x'}$ are isomorphic, choose any column in T_x and similarly for $T_{x'}$. We claim that $\varphi(m, n) = (m, n + km)$ is an isomorphism between T_x and $T_{x'}$.

To see this, note that φ sends m to m and n to $m + kn$, both of which are linear, so

we certainly have that φ is bijective. Further, we notice that adjacency of vertices is preserved as, if $(m, n) \rightarrow (m', n')$ in T_x then we claim that $(m, n+km) \rightarrow (m', n'+km')$ in T'_x . To check this, we refer to the above cases where $(m, n) \rightarrow (m', n')$.

If $m = m'$ and $n > n'$ then $\varphi(m, n) \rightarrow \varphi(m', n')$ as $m = m'$ and so $n + km > n' + km'$. If $m + 1 = m'$ and $x(n' - n) = 1$ then we arrive at the same conclusion as $m + 1 = m'$ and $x'((n' + km') - (n + km)) = x'(n' - n + k) = x(n' - n) = 1$. Finally, if $|m - m'| \geq 2$ and $m < m'$ then it still remains that $\varphi(m, n) \rightarrow \varphi(m', n')$ as φ leaves m and m' unchanged. Thus, we are able to conclude that $\varphi : T_x \rightarrow T'_x$ is an isomorphism.

For the other direction, consider $T_x \cong T_{x'}$. That is, $\varphi : T_x \rightarrow T_{x'}$ gives an isomorphism as described. To see that $x \in E_{\mathbb{Z}}x'$, we must first see that we can recover x from T_x . In order to do so, we require that x have sufficiently many 0 and 1 values. To be precise, we require at least one of the following conditions:

- (i) For every n there exists some $k < n$ such that $x(k - n) = 1$ and $x(k) = 0$
- (ii) For every n there exists some k such that $x(-k) = 0$ and $x(k - n) = 0$

Note that, excluding the trivial case where $w = (-1, n)$, each of these conditions on x are G_δ and dense, hence comeager. With these requirements, we now come to the definition of the comeager set \mathcal{C} containing all $x \in 2^{\mathbb{Z}}$ satisfying the above conditions.

By the above conditions, for any vertex v , we can identify the five columns surrounding v which we will denote as S_v . More specifically, S_v is the set of points involved in a three-cycle with v .

To see this, let $v = (0, 0)$ and consider a second vertex w . In the case that $w = (0, n)$, $n < 0$ is in the same column as v then condition (i) ensures that there is a third vertex u such that $w \rightarrow u$ and $u \rightarrow v$. Condition (i) ensures the same in

the case that $w = (-1, n)$ or $w = (1, n)$, $n \in \mathbb{Z}$. On the other hand, if $w = (-2, n)$ or $w = (2, n)$ then condition (ii) ensures that there exists a third vertex u such that $w \rightarrow u$ and $u \rightarrow v$.

We note that it is impossible for there to be a three-cycle involving two vertices that are more than two columns apart. To see that this is the case, suppose that $v = (0, 0)$ and $w = (m, n)$ for $|m| \geq 3$ and for some $n \in \mathbb{Z}$. Further, suppose there is a third vertex u such that $w \rightarrow u$. If $m \leq -3$ then $w \rightarrow v$ and, by assumption $w \rightarrow u$. Therefore w cannot be involved in a three-cycle with v . So now, if we consider $m \geq 3$, we certainly have $v \rightarrow w$. Since $w \rightarrow u$ then it must be that $m' \geq m - 1$ and so $m' \geq 2$. That is, $|0 - m'| = |m'| \geq 2$ and so $v \rightarrow u$. Once again, we see that w cannot be involved in a three-cycle with v based on our construction.

Further, we can determine distinct columns within S_v as follows:

- (i) Let C_v denote the set of all vertices $w \in S_v$ such that $S_w = S_v$
- (ii) Let $C_{-2,v}$ denote the set of all vertices $w \in S_v$ such that v does not arrow anyone in C_w
- (iii) Let $C_{2,v}$ denote the set of all vertices $w \in S_v$ such that v is in $C_{-2,w}$
- (iv) Let $C_{-1,v}$ denote the set of all vertices $w \in S_v$ such that $w \notin C_{-2,v}$ and every $z \in C_{2,v}$ is contained in S_w
- (v) Let $C_{1,v}$ denote the set of all vertices $w \in S_v$ such that for every $z \in C_{-1,v}$, $w \in C_{2,z}$.

Focusing on $C_{1,v}$, we can recover the sequence x by choosing any v and looking at edges between v and each $w \in C_{1,v}$. If we repeat this process for $T_{x'}$ we then

have x and x' . Well, as φ is an isomorphism between T_x and $T_{x'}$, φ must map $C_{1,v}$ (a \mathbb{Z} -ordered subgraph) in T_x to $C_{1,\varphi(v)}$ in $T_{x'}$ in an order preserving way. The only possible map is a shift. That is, the edges from v to vertices in $C_{1,v}$ in T_x are a shift of the edges from $\varphi(v)$ to vertices in $C_{1,\varphi(v)}$ in $T_{x'}$. Well, as x and x' were used to define these edges respectively it must be the case that x is a shift of x' and so $x E_{\mathbb{Z}} x'$ as expected. \square

Finally, we note that the \cong_{VTT} is strictly more complex than E_0 . That is, we recall the existence of a Δ_2^1 reduction from E_{ω_1} to the isomorphism relation on vertex-transitive linear orders. As linear orders are tournaments we also get that there is a Δ_2^1 reduction from E_{ω_1} to the isomorphism relation on vertex-transitive tournaments. If it were the case that $E_0 \sim_B \cong_{VTT}$ then there would exist a Borel reduction from \cong_{VTT} to E_0 . But that would mean E_{ω_1} was reducible to E_0 which cannot be the case as E_0 and E_{ω_1} are incomparable (refer to the remark following Corollary 3.3 in [7]) and so we reach a contradiction.

REFERENCES

- [1] John D. Clemens. Isomorphism of homogeneous structures. *Notre Dame Journal of Formal Logic*, 50(1):1–22, 2009.
- [2] R. Dougherty, S. Jackson, and A. S. Kechris. The structure of hyperfinite Borel equivalence relations. *Trans. Amer. Math. Soc.*, 341(1):193–225, 1994.
- [3] Harvey Friedman and Lee Stanley. A Borel reducibility theory for classes of countable structures. *J. Symbolic Logic*, 54(3):894–914, 1989.
- [4] Su Gao. *Invariant descriptive set theory*, volume 293 of *Pure and Applied Mathematics (Boca Raton)*. CRC Press, Boca Raton, FL, 2009.
- [5] Joel David Hamkins and Andy Lewis. Infinite time Turing machines. *J. Symbolic Logic*, 65(2):567–604, 2000.
- [6] L. A. Harrington, A. S. Kechris, and A. Louveau. A Glimm-Effros dichotomy for Borel equivalence relations. *J. Amer. Math. Soc.*, 3(4):903–928, 1990.
- [7] Greg Hjorth. An absoluteness principle for Borel sets. *J. Symbolic Logic*, 63(2):663–693, 1998.
- [8] Alan H. Mekler. Stability of nilpotent groups of class 2 and prime exponent. *J. Symbolic Logic*, 46(4):781–788, 1981.

- [9] Joseph G. Rosenstein. *Linear orderings*, volume 98 of *Pure and Applied Mathematics*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1982.